

## Minocycline May be Useful to Prevent or Treat Methamphetamine-Induced Neural Cell Death: Hypothetic Role of Autophagia and Apoptosis Signaling Pathway

Methamphetamine (METH) is a psychostimulant agent, and its neurotoxic and neurobehavioral consequences are serious.<sup>[1]</sup> Studies indicated that chronic abuse of METH can cause neural cell death in human and animal subjects.<sup>[1,2]</sup> Previous studies also showed that chronic abuse of METH can cause neurodegeneration, clear mechanism to these types of neurodegeneration remains unclear, but it seems that neuroinflammatory signaling pathways involved in this manner.<sup>[1]</sup> By induction of Bcl-2 phosphorylation after beginning an autophagy or apoptosis signal on the cell surface, in particular by activating tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) receptor, c-Jun N-terminal kinase (JNK) causes phosphorylation of Bcl-2 which this phenomenon causes inactivation and dissociation of Bcl-2 from Beclin1 or Bax which cause initiation of autophagy or apoptosis respectively and therefore cell death was

occurred<sup>[3,4]</sup> On the other way, during recent years, using new neuroprotective compounds with therapeutic probability for the treatment of drug abuses induced sequels has been amazingly increased.<sup>[5,6]</sup> Minocycline is a broad-spectrum and long-acting antibiotic and possesses anti-inflammatory, neuroprotective, and neural survival properties.<sup>[7,8]</sup> Pervious data suggested that minocycline has anti-inflammatory properties and can inhibit TNF- $\alpha$  receptor downstream signaling cascade in the neural cells.<sup>[9,10]</sup> Thus, according to the above evidence, we can suggest and have the theory that by inhibition of TNF-receptor activation and downstream JNK/Bcl-2-Beclin1 or Bcl-2/Bax signaling pathways, minocycline will likely inhibit METH-induced apoptosis and autophagia and probably by this mechanism can inhibit METH-induced neurodegeneration and neural cell loss [Figure 1].




Figure 1: Minocycline can inhibit METH-induced TNFR activation and c-Jun N-terminal kinase/Bcl-2-Beclin1 or c-Jun N-terminal kinase/Bcl-2/Bax signaling pathway which may reduce METH-induced autophagia and apoptosis and consequences neurodegeneration. Further, minocycline directly activates mitochondrial biogenesis which leads to inhibition of METH-induced oxidative stress and inflammation. TNFR: TNF- $\alpha$  receptor, METJ: Methamphetamine

**Afrah Sepehr, Majid Motaghinejad, Soomaayeh Heysieattalab<sup>1</sup>, Sepideh Safari**

Razi Drug Research Center, Iran University of Medical Sciences, Tehran,  
<sup>1</sup>Division of Cognitive Neuroscience, University of Tabriz, Tabriz, Iran

**Address for correspondence:**

Dr. Majid Motaghinejad,  
Tehran, Hemmat Highway Beside the Milad Tower, Iran University of  
Medical Sciences, P.O. Box: 1449614535. Tehran, Iran.  
E-mail: [Motaghinezhad.m@iums.ac.ir](mailto:Motaghinezhad.m@iums.ac.ir)

## References

1. Rahimi Borumand M, Motaghinejad M, Motevalian M, Gholami M. Duloxetine by modulating the Akt/GSK3 signaling pathways has neuroprotective effects against methamphetamine-induced neurodegeneration and cognition impairment in rats. *Iran J Med Sci* 2019; 44:146-54.
2. Casaletto KB, Obermeit L, Morgan EE, Weber E, Franklin DR, Grant I, *et al.* Depression and executive dysfunction contribute to a metamemory deficit among individuals with methamphetamine use disorders. *Addict Behav* 2015; 40:45-50.
3. Olmos G, Lladó J. Tumor necrosis factor alpha: A link between neuroinflammation and excitotoxicity. *Mediators Inflamm* 2014;2014: 861231
4. Poon HF, Abdullah L, Mullan MA, Mullan MJ, Crawford FC. Cocaine-induced oxidative stress precedes cell death in human neuronal progenitor cells. *Neurochem Int* 2007; 50:69-73.
5. Elkashef A, Vocci F, Hanson G, White J, Wickes W, Tiihonen J. Pharmacotherapy of methamphetamine addiction: An update. *Subst Abus* 2008; 29:31-49.
6. Ballester J, Valentine G, Sofuoglu M. Pharmacological treatments for methamphetamine addiction: Current status and future directions. *Expert Rev Clin Pharmacol* 2017; 10:305-14.
7. Miao H, Li R, Han C, Lu X, Zhang H. Minocycline promotes posthemorrhagic neurogenesis via M2 microglia polarization via upregulation of the TrkB/BDNF pathway in rats. *J Neurophysiol* 2018; 120:1307-17.
8. Scott G, Zetterberg H, Jolly A, Cole JH, De Simoni S, Jenkins PO, *et al.* Minocycline reduces chronic microglial activation after brain trauma but increases neurodegeneration. *Brain* 2018; 141:459-71.
9. Stirling DP, Koochesfahani KM, Steeves JD, Tetzlaff W. Minocycline as a neuroprotective agent. *Neuroscientist* 2005; 11:308-22.
10. Lee SM, Yune TY, Kim SJ, Kim YC, Oh YJ, Markelonis GJ, *et al.* Minocycline inhibits apoptotic cell death via attenuation of TNF-alpha expression following iNOS/NO induction by lipopolysaccharide in neuron/glia co-cultures. *J Neurochem* 2004; 91:568-78.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online	
<b>Quick Response Code:</b>	<b>Website:</b>
	<a href="http://www.advbiores.net">www.advbiores.net</a>
	<b>DOI:</b>
	10.4103/abr.abr_258_19

**How to cite this article:** Sepehr A, Motaghinejad M, Heysieattalab S, Safari S. Minocycline may be useful to prevent or treat methamphetamine-induced neural cell death: Hypothetic role of autophagia and apoptosis signaling pathway. *Adv Biomed Res* 2020;9:7.

**Received:** 12 October 2019; **Revised:** 10 December 2019;  
**Accepted:** 16 December 2019; **Published:** 25 February 2020