

The Interaction of *Helicobacter pylori* Infection and Type 2 Diabetes Mellitus

Abstract

Helicobacter pylori is one of the most common human pathogens that can cause gastrointestinal (GI) disorders, including simple gastritis, gastric ulcer, and malignant gastritis. In some cases, such as immunodeficiency and underlying diseases, it can be problematic as opportunistic infections. Diabetes mellitus (type 2) (T2DM) is one of the *H. pylori* underlying diseases. Since GI problems are observed in diabetic patients, it is necessary to treat *H. pylori* infection. In this review, we aimed to evaluate the possible relationship between *H. pylori* and T2DM according to epidemiological surveys of 70 studies retrieved from databases, including Scopus, PubMed, and Google Scholar about the relationship between *H. pylori* and T2DM, and discuss the reported background mechanisms of this correlation. According to the results of our study, the different studies have shown that *H. pylori* is more prevalent in Type 2 diabetic patients than healthy individuals or nondiabetic patients. The reason is development of *H. pylori* infection-induced inflammation and production of inflammatory cytokines as well as different hormonal imbalance by this bacterium, which are associated with diabetes mellitus. On the other hand, by tracing anti-*H. pylori* antibodies in patients with diabetes mellitus and occurrence of symptoms such as digestive problems in >75% of these patients, it can be concluded that there is a relationship between this bacterium and T2DM. Considering the evidence, it is crucially important that the probability of infection with *H. pylori* is evaluated in patients with T2DM so that medical process of the patient is followed with higher cautious.

Keywords: Diabetes, diabetes mellitus, *Helicobacter pylori*, Type 2 diabetes

Introduction

In 1983, Warren (a biologist) and Marshall (a clinician) described *Helicobacter pylori*.^[1] First, they named the bacterium *Campylobacter pyloridis*, and then, it was named *Campylobacter pylori*. Diagnosis and treatment of the upper gastroduodenal disease have been changed dramatically since 20 years ago that *H. pylori* was cultured for the first time.^[2-4] Peptic ulcer disease is now approached as an infectious disease.^[5] The role of *H. pylori* infection is increasingly recognized in gastric cancers as well as evaluating its role in other gastrointestinal (GI) diseases.^[6]

Elevated antibodies level against *H. pylori* also attracted the attention to some extra-gastric diseases, including diabetes mellitus.^[7,8]

Among the patients referring to diabetes clinics, as many as 75% of them will report significant GI symptoms.^[9,10] The

entire GI tract can be affected by diabetes from the oral cavity and esophagus to the large bowel and anorectal region. Thus, the experienced symptom complex may vary widely.^[11] The common complaints can include dysphagia, early satiety, reflux, constipation, abdominal pain, nausea, vomiting, and diarrhea. Many patients remain undiagnosed and undertreated because the GI tract has not been conventionally associated with diabetes and its complications.^[12]

Type 2 diabetes mellitus (T2DM) is turning to be pandemic so that it is responsible for death of 3.8 million of the adult population in the world and regarded as a serious risk for public health.^[13,14] Increasing blood sugar, which is chronically observed in the patients with diabetes, may cause long-term damage to different organs, especially eyes, kidney, nervous system, heart, and blood vessels. At least 80% of the patients with diabetes will die due to consequences of cardiac complications.^[14-16]

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

**Seyed Abolfazi
Hosseininasab
Nodoushan,
Amin Nabavi¹**

From the Departments of Microbiology and ¹Biochemistry, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Address for correspondence:

Mr. Seyed Abolfazi
Hosseininasab Nodoushan,
Department of Microbiology,
School of Medicine, Isfahan
University of Medical
Sciences, Isfahan, Iran.
E-mail: majidhoseini1001@yahoo.com

Access this article online

Website: www.advbiores.net

DOI: 10.4103/abr.abr_37_18

Quick Response Code:



How to cite this article: Hosseininasab Nodoushan SA, Nabavi A. The Interaction of *Helicobacter pylori* Infection and Type 2 Diabetes Mellitus. *Adv Biomed Res* 2019;8:15.

Received: March, 2018. **Accepted:** January, 2019.

Pathogenic mechanisms of diabetes mellitus include insulin resistance (IR), chronic inflammation, insufficiency of insulin secretion (due to impaired pancreatic beta-cells), glucose toxicity, and lipotoxicity.^[14]

Relationship between *Helicobacter pylori* and Type 2 Diabetes Mellitus

Evidence indicates that diabetes may accompany by *H. pylori* infection, which chronic and insulin-resistant inflammation may increase the risk for T2DM. In addition, gastritis resulting from *H. pylori* may potentially affect gut-related hormones and inflammatory cytokines.^[14,17]

Although there is no strong evidence for this relationship, some reasons can be considered to discuss it, which are summarized in the following:

First, diabetes causes impairment in the function of the cellular and humoral immunity, which also increases the individual's sensitivity to *H. pylori* infection.^[18] Second, it reduces GI movements and secretion of gastric acid, which in turn increases colonization and bacterial infections.^[19] Third, changes in glucose metabolism may alter chemical production in the gastric mucosa, which results in colonization of more bacteria.^[20] Ultimately, diabetic patients are more likely to be exposed to pathogens than healthy people, due to their more presence in the hospital environment.^[21]

There is controversy about the link between *H. pylori* infection and diabetes as some studies indicate a higher prevalence of infection in diabetic patients,^[22-24] whereas in the others, no difference has been reported.^[25-27] Jeon *et al.* have been reported for the first time that *H. pylori* infection leads to increase the incidence of T2DM using a prospective cohort of 782 Latino individuals older than 60 years.^[19] This study showed that people with *H. pylori* infection would more suffer from diabetes in comparison to healthy individuals.

It is found that 84.6% of diabetic patients with *H. pylori* infection had diabetes for >10 years. Besides the glycemic control, diabetes duration is the main risk factor of increasing the risk of chronic diabetes-related complications, which its importance in our study is the autonomic neuropathy and gastropathy that are critical predictors for *H. pylori* infection in diabetics.^[12,28]

Bayati *et al.* in a study concluded that 61.5% of those with *H. pylori*-positive status had one or more of the chronic diabetic complications. This finding is in agreement with the findings of a study indicating that diabetic patients with a history of retinopathy, nephropathy, or neuropathy should be presupposed to have GI abnormalities until proven otherwise.^[29]

Role of Inflammatory Cytokines

At the initial phases of infection, polymorphonuclear cells penetrate into GI mucus. After a while, these cells

are replaced by mononuclear cells at chronic phase of infection. Mononuclear cells are specified by the production of inflammatory cytokines, which in addition to local effect can be effective also as released on other tissues and organs. It causes increase in the extraintestinal diseases in people with *H. pylori* infection, such as cardiovascular, neurological, autoimmune, thyroid, liver, and biliary diseases.^[30] Therefore, the bacteria cause to develop inflammation and production of different cytokines and impairment in absorption of nutrients and medicines and can lead to the induction and development of various diseases.^[31-35] In a study by Zojaji *et al.* on 85 patients referring to a endocrinology clinic of Shahid Beheshti University of Medical Sciences in Tehran, Iran, it concluded that the most important produced inflammatory cytokines include C-reactive protein (CRP), interleukin-6 (IL-6), and tumor necrosis factor alpha (TNF- α), which contribute to IR and diabetes incidence.^[36-41] High levels of CRP (high-sensitive CRP [hsCRP]) are important determinants in diabetes-related research so that the most papers reported strong relationship between CRP level and diabetes risk. Various studies introduced this factor as the connection between *H. pylori* and Type 2 diabetes; as of 11 prospective studies, seven reported a significant positive association between hsCRP levels and diabetes risk^[30,42-48] while it found no association in the others.^[49-52] Various studies have identified this factor as a bridge between *H. pylori* and Type 2 diabetes as El Hadidy *et al.* reported a significant increase in CRP level and *H. pylori* infection among Type 2 diabetic patients as well as Oshima *et al.* found a significant increase in CRP level in nonsmoker healthy individuals seropositive for *H. pylori* infection. Markus *et al.* found significant association between CRP and *H. pylori* seropositivity.^[30,37,53]

A study by Aydemir *et al.* showed higher homeostatic model assessment-estimated IR scores in *H. pylori*-positive individuals.^[54] Furthermore, a Japanese study in 2009 conducted on a large population of 1107 asymptomatic individuals showed that *H. pylori* was significantly and independently related to IR.^[55] However, Gillum supported that there are no consistent associations between *H. pylori* infection and prevalence of diabetes or syndrome-related variables of the IR in American men of 40–74 years.^[56] Furthermore, Park *et al.* reported that metabolic and inflammatory parameters, including blood sugar, lipid profiles, IR, white blood cell count, and CRP levels, were not changed after *H. pylori* eradication. It is worth noting that *H. pylori* infection was not determined in all studies by organisms' histologic detection of mucosal biopsy specimens, which is considered the diagnostic gold standard.^[14,57]

Role of Hormones

Helicobacter by induction of gastritis is potentially able to influence the secretion of hormones, such as leptin, ghrelin,

gastrin, and somatostatin, and subsequently increases the diabetes risk. Gastrin increases insulin secretion and somatostatin leads to reduction. *Helicobacter* causes to increase gastrin concentration and lower serum somatostatin concentration through influencing these hormones and thus affects insulin secretion.^[58-60]

Reduced insulin secretion is one of the major factors in the incidence of Type 2 diabetes. In the studies by So *et al.*, it indicated that *H. pylori* interdependently can predict dysfunction of pancreatic beta-cells in Chinese males.^[61]

In addition, Rahman *et al.* found a positive relationship between *H. pylori* infection and defective insulin secretion so that insulin molecules are highly susceptible to damage from oxidative stress and inflammation. Hence, *Helicobacter* inflammation probably causes defection in insulin production.^[62]

In addition, in the study by Hsieh *et al.*, it reported that patients infected by *H. pylori*, especially at young ages are defective in insulin secretion, which is considered as a risk factor for Type 2 diabetes.^[63] In addition, Gen *et al.* proved that elimination of pylori infection considerably reduces insulin level.^[64]

Ghrelin causes reduction of energy consumption and increasing the weight, whereas leptin reduces food absorption, on the one hand, and increases energy consumption, on the other hand.^[60,65]

It has been shown that *H. pylori* infection impairs ghrelin production and enhances leptin production.^[66-68] Low ghrelin levels are associated with elevated fasting insulin concentrations, IR, and T2DM.^[69] Leptin has also been implicated in IR development, and elevated levels correlate with IR in lean men and patients with T2DM^[70-72] [Figure 1].

Insulin resistance and abnormal insulin secretion are central to the development of Type 2 diabetes mellitus. On the one

hand, *Helicobacter pylori* infection brings about chronic low-grade inflammation with upregulation of several cytokines such as C-reactive protein, tumor necrosis factor, and interleukin-1 β , which may impact insulin action and pancreatic β -cell secretion. On the other hand, *Helicobacter pylori*-induced gastritis can potentially affect gastric hormones secretion, including leptin, ghrelin, gastrin, and somatostatin, which could affect insulin sensitivity and glucose homeostasis. In addition, other mechanisms and mediators may be involved in the possible causative relationship between *Helicobacter pylori* infection and Type 2 diabetes mellitus.^[73,74]

Helicobacteria Can also Be Associated with Other Diseases due to Type 2 Diabetes

Diabetic patients are at risk of cardiovascular and thrombo-occlusive cerebral diseases. There is a possible correlation between *H. pylori* infection and cardiovascular or cerebrovascular diseases. Chronic *H. pylori* colonization may be in association with an increased risk of atherosclerosis. It is hypothesized that *H. pylori* infection has atherogenic capacities by activating chronic low grade of homeostasis cascade.^[73-75] The association between *H. pylori* and acute cerebrovascular disease may be due to a higher prevalence of virulent *H. pylori* strains in atherosclerotic stroke patients.^[76]

El Hadidy *et al.* also found a significant increase in fibrinogen level as one of the risk factors of atherosclerosis in *H. pylori*-positive diabetic patients in comparison to negative patients.^[30] The same results were found by Schumacher *et al.* in patients with coronary heart disease.^[77] Zito *et al.* reported the increased level of plasma fibrinogen in *H. pylori*-infected patients even after controlling of possible related confounding factors either infection or fibrinogen.^[78]

Conclusions

In this regard by studying various studies, we can conclude that:

1. *H. pylori* infection is more common in diabetic patients
2. There is higher frequency of *H. pylori* infection among elderly diabetics
3. *H. pylori* infection enhances the presence of chronic diabetic complications in its different categories
4. There is no significant difference in male-to-female ratio distribution of *H. pylori* infection.

However, as already mentioned, there are opposite results in this regard, though diabetes mellitus is a multifaceted and multistep disease that is improbably resulted from a single cause, as the risk factors that deserve attention include GI infections and the intestinal microbiota composition. Evidence supporting an etiological role of *H. pylori* in T2DM developing would indicate that preventive measures, such as improving hygiene and treatments using




Figure 1: Potential mechanisms for *Helicobacter pylori* contribution to Type 2 diabetes mellitus

antibiotics and proton-pump inhibitor combinations, should be explored as the target of interventions in high-risk communities.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

1. Marshall BJ, Warren JR. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet* 1984;1:1311-5.
2. Papamichael KX, Papaioannou G, Karga H, Roussos A, Mantzaris GJ. *Helicobacter pylori* infection and endocrine disorders: Is there a link? *World J Gastroenterol* 2009;15:2701-7.
3. Suerbaum S, Michetti P. *Helicobacter pylori* infection. *N Engl J Med* 2002;347:1175-86.
4. Khademi F, Faghri J, Poursina F, Esfahani BN, Moghim S, Fazeli H, et al. Resistance pattern of *Helicobacter pylori* strains to clarithromycin, metronidazole, and amoxicillin in Isfahan, Iran. *J Res Med Sci* 2013;18:1056-60.
5. Malfertheiner P, Chan FK, McColl KE. Peptic ulcer disease. *Lancet* 2009;374:1449-61.
6. Hosseini E, Poursina F, de Wiele TV, Safaei HG, Adibi P. *Helicobacter pylori* in Iran: A systematic review on the association of genotypes and gastroduodenal diseases. *J Res Med Sci* 2012;17:280-92.
7. Feldman R. Epidemiologic observations and open questions about disease and infection caused by *Helicobacter pylori*. In: Achtman M, Serbaum S, editors. *Helicobacter pylori: Molecular and Cellular Biology*. Uk: Horizon Scientific, 2001:29-51.
8. Malaty HM, Graham DY. Importance of childhood socioeconomic status on the current prevalence of *Helicobacter pylori* infection. *Gut* 1994;35:742-5.
9. Frenck RW Jr., Clemens J. *Helicobacter* in the developing world. *Microbes Infect* 2003;5:705-13.
10. Granquist A, Bredberg A, Sveger T, Axelsson I. A longitudinal cohort study on the prevalence of *Helicobacter pylori* antibodies in Swedish children and adolescents. *Acta Paediatr* 2002;91:636-40.
11. Rajashekhar V, Bhasin DK, Ray P, Vaiphei K, Sharma BC, Singh K, et al. *Helicobacter pylori* infection in chronic smokers with non ulcer dyspepsia. *Trop Gastroenterol* 2000;21:71-2.
12. Bakka AS, Salih BA. Prevalence of *Helicobacter pylori* infection in asymptomatic subjects in Libya. *Diagn Microbiol Infect Dis* 2002;43:265-8.
13. van Dieren S, Beulens JW, van der Schouw YT, Grobbee DE, Neal B. The global burden of diabetes and its complications: An emerging pandemic. *Eur J Cardiovasc Prev Rehabil* 2010;17 Suppl 1:S3-8.
14. He C, Yang Z, Lu NH. *Helicobacter pylori* infection and diabetes: Is it a myth or fact? *World J Gastroenterol* 2014;20:4607-17.
15. Agardh E, Allebeck P, Hallqvist J, Moradi T, Sidorchuk A. Type 2 diabetes incidence and socio-economic position: A systematic review and meta-analysis. *Int J Epidemiol* 2011;40:804-18.
16. Qi L, Hu FB, Hu G. Genes, environment, and interactions in prevention of Type 2 diabetes: A focus on physical activity and lifestyle changes. *Curr Mol Med* 2008;8:519-32.
17. Chen Y, Blaser MJ. Association between gastric *Helicobacter pylori* colonization and glycated hemoglobin levels. *J Infect Dis* 2012;205:1195-202.
18. Borody T, Ren Z, Pang G, Clancy R. Impaired host immunity contributes to *Helicobacter pylori* eradication failure. *Am J Gastroenterol* 2002;97:3032-7.
19. Jeon CY, Haan MN, Cheng C, Clayton ER, Mayeda ER, Miller JW, et al. *Helicobacter pylori* infection is associated with an increased rate of diabetes. *Diabetes Care* 2012;35:520-5.
20. de Luis DA, de la Calle H, Roy G, de Argila CM, Valdezate S, Canton R, et al. *Helicobacter pylori* infection and insulin-dependent diabetes mellitus. *Diabetes Res Clin Pract* 1998;39:143-6.
21. Gentile S, Turco S, Oliviero B, Torella R. The role of autonomic neuropathy as a risk factor of *Helicobacter pylori* infection in dyspeptic patients with Type 2 diabetes mellitus. *Diabetes Res Clin Pract* 1998;42:41-8.
22. Devrajani BR, Shah SZ, Soomro AA, Devrajani T. Type 2 diabetes mellitus: A risk factor for *Helicobacter pylori* infection: A hospital based case-control study. *Int J Diabetes Dev Ctries* 2010;30:22-6.
23. Bener A, Micallef R, Afifi M, Derbala M, Al-Mulla HM, Usmani MA, et al. Association between Type 2 diabetes mellitus and *Helicobacter pylori* infection. *Turk J Gastroenterol* 2007;18:225-9.
24. Gulcelik NE, Kaya E, Demirbas B, Culha C, Koc G, Ozkaya M, et al. *Helicobacter pylori* prevalence in diabetic patients and its relationship with dyspepsia and autonomic neuropathy. *J Endocrinol Invest* 2005;28:214-7.
25. Anastasios R, Goritsas C, Papamihail C, Trigidou R, Garzonis P, Ferti A, et al. *Helicobacter pylori* infection in diabetic patients: Prevalence and endoscopic findings. *Eur J Intern Med* 2002;13:376.
26. Ko GT, Chan FK, Chan WB, Sung JJ, Tsoi CL, To KF, et al. *Helicobacter pylori* infection in Chinese subjects with Type 2 diabetes. *Endocr Res* 2001;27:171-7.
27. Stanciu OG, Trifan A, Sfarti C, Cojocariu C, Stanciu C. *Helicobacter pylori* infection in patients with diabetes mellitus. *Rev Med Chir Soc Med Nat Iasi* 2003;107:59-65.
28. Parsonnet J, Shmueli H, Haggerty T. Fecal and oral shedding of *Helicobacter pylori* from healthy infected adults. *JAMA* 1999;282:2240-5.
29. Kozák R, Juhász E, Horvát G, Harsa E, Lövei L, Sike R, et al. *Helicobacter pylori* infection in diabetic patients. *Orv Hetil* 1999;140:993-5.
30. El Hadidy M, Abdul-Aziz MY, Mokhtar AR, El Ata MM, El Gwad SS. *Helicobacter pylori* infection and vascular complications in patients with Type 2 diabetes mellitus. *J Taibah Univ Med Sci* 2009;4:62-72.
31. Shoelson SE, Lee J, Goldfine AB. Inflammation and insulin resistance. *J Clin Invest* 2006;116:1793-801.
32. Fernández-Real JM, López-Bermejo A, Vendrell J, Ferri MJ, Recasens M, Ricart W, et al. Burden of infection and insulin resistance in healthy middle-aged men. *Diabetes Care* 2006;29:1058-64.
33. Howard BV, Best L, Comuzzie A, Ebbesson SO, Epstein SE, Fabsitz RR, et al. C-reactive protein, insulin resistance, and metabolic syndrome in a population with a high burden of subclinical infection: Insights from the genetics of coronary artery disease in Alaska natives (GOCADAN) study. *Diabetes Care* 2008;31:2312-4.
34. Misiewicz JJ. Current insights in the pathogenesis of *Helicobacter pylori* infection. *Eur J Gastroenterol Hepatol* 1995;7:701-3.

35. Mishra RR, Tewari M, Shukla HS. Association of *Helicobacter pylori* infection with inflammatory cytokine expression in patients with gallbladder cancer. *Indian J Gastroenterol* 2013;32:232-5.
36. Diomedì M, Stanzione P, Sallustio F, Leone G, Renna A, Misaggi G, et al. Cytotoxin-associated gene-A-positive *Helicobacter pylori* strains infection increases the risk of recurrent atherosclerotic stroke. *Helicobacter* 2008;13:525-31.
37. Oshima T, Ozono R, Yano Y, Oishi Y, Teragawa H, Higashi Y, et al. Association of *Helicobacter pylori* infection with systemic inflammation and endothelial dysfunction in healthy male subjects. *J Am Coll Cardiol* 2005;45:1219-22.
38. Jackson L, Britton J, Lewis SA, McKeever TM, Atherton J, Fullerton D, et al. A population-based epidemiologic study of *Helicobacter pylori* infection and its association with systemic inflammation. *Helicobacter* 2009;14:108-13.
39. Hamed SA, Amine NF, Galal GM, Helal SR, Tag El-Din LM, Shawky OA, et al. Vascular risks and complications in diabetes mellitus: The role of *Helicobacter pylori* infection. *J Stroke Cerebrovasc Dis* 2008;17:86-94.
40. Wellen KE, Hotamisligil GS. Inflammation, stress, and diabetes. *J Clin Invest* 2005;115:1111-9.
41. Zojaji H, Ataei E, Sherafat SJ, Ghobakhlou M, Fatemi SR. The effect of the treatment of *Helicobacter pylori* infection on the glycemic control in Type 2 diabetes mellitus. *Gastroenterol Hepatol Bed Bench* 2013;6:36-40.
42. Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM. C-reactive protein, interleukin 6, and risk of developing Type 2 diabetes mellitus. *JAMA* 2001;286:327-34.
43. Barzilay JI, Abraham L, Heckbert SR, Cushman M, Kuller LH, Resnick HE, et al. The relation of markers of inflammation to the development of glucose disorders in the elderly: The cardiovascular health study. *Diabetes* 2001;50:2384-9.
44. Duncan BB, Schmidt MI, Pankow JS, Ballantyne CM, Couper D, Vigo A, et al. Low-grade systemic inflammation and the development of Type 2 diabetes: The atherosclerosis risk in communities study. *Diabetes* 2003;52:1799-805.
45. Freeman DJ, Norrie J, Caslake MJ, Gaw A, Ford I, Lowe GD, et al. C-reactive protein is an independent predictor of risk for the development of diabetes in the west of Scotland coronary prevention study. *Diabetes* 2002;51:1596-600.
46. Hu FB, Meigs JB, Li TY, Rifai N, Manson JE. Inflammatory markers and risk of developing Type 2 diabetes in women. *Diabetes* 2004;53:693-700.
47. Thorand B, Löwel H, Schneider A, Kolb H, Meisinger C, Fröhlich M, et al. C-reactive protein as a predictor for incident diabetes mellitus among middle-aged men: Results from the MONICA Augsburg Cohort Study, 1984-1998. *Arch Intern Med* 2003;163:93-9.
48. Laaksonen DE, Niskanen L, Nyssönen K, Punnonen K, Tuomainen TP, Valkonen VP, et al. C-reactive protein and the development of the metabolic syndrome and diabetes in middle-aged men. *Diabetologia* 2004;47:1403-10.
49. Festa A, D'Agostino R Jr, Tracy RP, Haffner SM; Insulin Resistance Atherosclerosis Study. Elevated levels of acute-phase proteins and plasminogen activator inhibitor-1 predict the development of Type 2 diabetes: The insulin resistance atherosclerosis study. *Diabetes* 2002;51:1131-7.
50. Han TS, Sattar N, Williams K, Gonzalez-Villalpando C, Lean ME, Haffner SM, et al. Prospective study of C-reactive protein in relation to the development of diabetes and metabolic syndrome in the Mexico City diabetes study. *Diabetes Care* 2002;25:2016-21.
51. Krakoff J, Funahashi T, Stehouwer CD, Schalkwijk CG, Tanaka S, Matsuzawa Y, et al. Inflammatory markers, adiponectin, and risk of Type 2 diabetes in the Pima Indian. *Diabetes Care* 2003;26:1745-51.
52. Snijder MB, Dekker JM, Visser M, Stehouwer CD, Yudkin JS, Bouter LM, et al. Prospective relation of C-reactive protein with Type 2 diabetes: Response to Han et al. *Diabetes Care* 2003;26:1656-7.
53. Markus HS, Risley P, Mendall MA, Steinmetz H, Sitzer M. *Helicobacter pylori* infection, the cytotoxin gene A strain, and carotid artery intima-media thickness. *J Cardiovasc Risk* 2002;9:1-6.
54. Aydemir S, Bayraktaroglu T, Sert M, Sokmen C, Atmaca H, Mungan G, et al. The effect of *Helicobacter pylori* on insulin resistance. *Dig Dis Sci* 2005;50:2090-3.
55. Gunji T, Matsuhashi N, Sato H, Fujibayashi K, Okumura M, Sasabe N, et al. *Helicobacter pylori* infection significantly increases insulin resistance in the asymptomatic Japanese population. *Helicobacter* 2009;14:144-50.
56. Gillum RF. Infection with *Helicobacter pylori*, coronary heart disease, cardiovascular risk factors, and systemic inflammation: The third national health and nutrition examination survey. *J Natl Med Assoc* 2004;96:1470-6.
57. Park SH, Jeon WK, Kim SH, Kim HJ, Park DI, Cho YK, et al. *Helicobacter pylori* eradication has no effect on metabolic and inflammatory parameters. *J Natl Med Assoc* 2005;97:508-13.
58. Jeffery PL, McGuckin MA, Linden SK. Endocrine impact of *Helicobacter pylori*: Focus on ghrelin and ghrelin o-acyltransferase. *World J Gastroenterol* 2011;17:1249-60.
59. Roper J, Francois F, Shue PL, Mourad MS, Pei Z, Olivares de Perez AZ, et al. Leptin and ghrelin in relation to *Helicobacter pylori* status in adult males. *J Clin Endocrinol Metab* 2008;93:2350-7.
60. Kaneko H, Konagaya T, Kusugami K. *Helicobacter pylori* and gut hormones. *J Gastroenterol* 2002;37:77-86.
61. So WY, Tong PC, Ko GT, Ma RC, Ozaki R, Kong AP, et al. Low plasma adiponectin level, white blood cell count and *Helicobacter pylori* titre independently predict abnormal pancreatic beta-cell function. *Diabetes Res Clin Pract* 2009;86:89-95.
62. Rahman MA, Cope MB, Sarker SA, Garvey WT, Chaudhury HS, Khaled MA, et al. *Helicobacter pylori* infection and inflammation: Implication for the pathophysiology of diabetes and coronary heart disease in Asian Indians. *J Life Sci* 2009;1:45-50.
63. Hsieh MC, Wang SS, Hsieh YT, Kuo FC, Soon MS, Wu DC, et al. *Helicobacter pylori* infection associated with high HbA1c and Type 2 diabetes. *Eur J Clin Invest* 2013;43:949-56.
64. Gen R, Demir M, Ataseven H. Effect of *Helicobacter pylori* eradication on insulin resistance, serum lipids and low-grade inflammation. *South Med J* 2010;103:190-6.
65. Calam J. *Helicobacter pylori* modulation of gastric acid. *Yale J Biol Med* 1999;72:195-202.
66. Osawa H, Nakazato M, Date Y, Kita H, Ohnishi H, Ueno H, et al. Impaired production of gastric ghrelin in chronic gastritis associated with *Helicobacter pylori*. *J Clin Endocrinol Metab* 2005;90:10-6.
67. Osawa H. Ghrelin and *Helicobacter pylori* infection. *World J Gastroenterol* 2008;14:6327-33.
68. Nishi Y, Isomoto H, Uotani S, Wen CY, Shikuwa S, Ohnita K, et al. Enhanced production of leptin in gastric fundic mucosa with *Helicobacter pylori* infection. *World J Gastroenterol* 2005;11:695-9.
69. Pöykkö SM, Kellokoski E, Hörkö S, Kauma H, Kesäniemi YA, Ukkola O, et al. Low plasma ghrelin is associated with insulin

- resistance, hypertension, and the prevalence of Type 2 diabetes. *Diabetes* 2003;52:2546-53.
70. Cohen B, Novick D, Rubinstein M. Modulation of insulin activities by leptin. *Science* 1996;274:1185-8.
 71. Segal KR, Landt M, Klein S. Relationship between insulin sensitivity and plasma leptin concentration in lean and obese men. *Diabetes* 1996;45:988-91.
 72. Fischer S, Hanefeld M, Haffner SM, Fusch C, Schwanebeck U, Köhler C, *et al.* Insulin-resistant patients with Type 2 diabetes mellitus have higher serum leptin levels independently of body fat mass. *Acta Diabetol* 2002;39:105-10.
 73. de Luis DA, Garcia Avello A, Lasuncion MA, Aller R, Martin de Argila C, Boixeda de Miquel D, *et al.* Improvement in lipid and haemostasis patterns after *Helicobacter pylori* infection eradication in Type 1 diabetic patients. *Clin Nutr* 1999;18:227-31.
 74. Danesh J, Whincup P, Walker M, Lennon L, Thomson A, Appleby P, *et al.* Low grade inflammation and coronary heart disease: Prospective study and updated meta-analyses. *BMJ* 2000;321:199-204.
 75. Sawayama Y, Ariyama I, Hamada M, Otaguro S, Machi T, Taira Y, *et al.* Association between chronic *Helicobacter pylori* infection and acute ischemic stroke: Fukuoka harasanshin atherosclerosis trial (FHAT). *Atherosclerosis* 2005;178:303-9.
 76. Pietroiusti A, Diomedi M, Silvestrini M, Cupini LM, Luzzi I, Gomez-Miguel MJ, *et al.* Cytotoxin-associated gene-A – Positive *Helicobacter pylori* strains are associated with atherosclerotic stroke. *Circulation* 2002;106:580-4.
 77. Schumacher A, Seljeflot I, Lerkerød AB, Sommervoll L, Otterstad JE, Arnesen H, *et al.* Positive chlamydia pneumoniae serology is associated with elevated levels of tumor necrosis factor alpha in patients with coronary heart disease. *Atherosclerosis* 2002;164:153-60.
 78. Zito F, Di Castelnuovo A, D'Orazio A, Negrini R, De Lucia D, Donati MB, *et al.* *Helicobacter pylori* infection and the risk of myocardial infarction: Role of fibrinogen and its genetic control. *Thromb Haemost* 1999;82:14-8.