

Immediate outcomes of eptifibatide therapy during intracoronary stent implantation

Hooman Shariati, Hamid Sanei, Ali Pourmoghadas, Leila Salehizadeh¹, Afshin Amirpour

Isfahan Cardiovascular Research Center, Cardiovascular Research Institute, ¹Interventional Cardiologist, Chamran Hospital, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract

Background: The objective of the present study was to assess the major immediate outcomes of eptifibatide therapy during intracoronary stent implantation.

Materials and Methods: In an interventional study, patients undergoing percutaneous coronary intervention (PCI) were randomized into either the eptifibatide ($n = 100$) or the control ($n = 107$) group. In each group, demographic and clinical characteristics such as cardiac death, stent thrombosis (ST), myocardial infarction (MI), rates of target lesion and vessel revascularization, cerebral vascular accident (CVA), and emergency coronary artery bypass grafting (CABG) were recorded.

Results: The overall rates of major adverse events such as mortality, Stent thrombosis (ST), Myocardial Infarction (MI), target lesion revascularization (TLR), target vessel revascularization (TVR), CVA, and emergency CABG within 24 h after stent implantation were low and comparable between the two groups; $P > 0.05$ considered significant for all comparisons.

Conclusion: There were no statistical differences between the clinical outcomes of groups administered with single-dose intracoronary eptifibatide and control groups among patients undergoing PCI during stent implantation.

Key Words: Eptifibatide, immediate clinical outcome, percutaneous coronary intervention, stent implantation

Address for correspondence:

Dr. Hamid Sanei, Isfahan Cardiovascular Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran.

E-mail: h_sanei@med.mui.ac.ir

Received: 25.06.2013, Accepted: 15.10.2014

INTRODUCTION

Coronary heart disease (CHD) is the narrowing or blockage of the coronary arteries generally caused by atherosclerosis.^[1] The outreach of stents has been a notable progress in the treatment of obstructive coronary artery disease (CAD) since the introduction of balloon angioplasty.^[2] Significant reduction in outcomes of mortality, myocardial infarction (MI), and target vessel

revascularization (TVR) using parenteral glycoprotein IIb/IIIa receptor antagonists has been widely reported for patients with acute coronary syndrome.^[3-6] Glycoprotein IIb/IIIa receptor antagonists consist of: The monoclonal antibody abciximab, the peptide receptor antagonist eptifibatide, and the nonpeptide receptor antagonists tirofiban and lamifiban.^[7] Among them eptifibatide (Integrilin) is often less expensive and more widely available in many hospitals.^[8,9]

Access this article online	
Quick Response Code:	Website: www.advbiores.net
	DOI: 10.4103/2277-9175.196831

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Shariati H, Sanei H, Pourmoghadas A, Salehizadeh L, Amirpour A. Immediate outcomes of eptifibatide therapy during intracoronary stent implantation. *Adv Biomed Res* 2016;5:204.

Eptifibatide has been shown to ameliorate cardiac outcomes in patients with percutaneous coronary intervention (PCI) by lessening the incidence of major adverse cardiac events.^[10-12] In an investigation, the complementary effect of thienopyridine pretreatment and the integrin blockade in coronary stent intervention were assessed; according to their results, pretreatment with thienopyridine decreased the rate of ischemic complications regardless of eptifibatide treatment. They expressed that the efficacy of eptifibatide is dependant on the presence of a loading dose of thienopyridine administration. Also, optimal outcomes were attained by receiving thienopyridine pretreatment along with eptifibatide therapy.^[13] In a study conducted between January 2003 and August 2005, the long-term mortality after bolus-only administration of abciximab, eptifibatide, and tirofiban during PCI were evaluated. As is shown in their results, eptifibatide was able to ameliorate long-term survival prognosis compared to abciximab when applied bolus-only.^[14] O'Shea *et al.* in a randomized controlled trial (RCT) entitled "Platelet glycoprotein IIb/IIIa Integrin blockade with Eptifibatide in coronary stent intervention" showed that adjunctive eptifibatide therapy during coronary stent implantation is beneficial within a 6-month follow-up.^[15] Intracoronary Eptifibatide bolus administration during PCI in patients with acute coronary syndromes indicated a higher local platelet glycoprotein IIb/IIIa receptor occupancy, which is related to improved microvascular perfusion demonstrated by improving the corrected thrombolysis in the MI frame count.^[16] In an investigation, different outcomes concerning the use of eptifibatide instead of abciximab in patients undergoing primary PCI were assessed. There were no differences expressed in the early outcomes of patients treated with eptifibatide, compared to patients treated with abciximab.^[8] Also, in another survey the clinical outcomes of eptifibatide and abciximab in acute ST elevation MI (STEMI) patients undergoing PCI were compared; no significant differences were observed in clinical outcomes between STEMI patients treated with eptifibatide or abciximab. It has been suggested that because of the lower cost of eptifibatide, it can be a suitable substitute for abciximab and reduce the overall medication costs while maintaining beneficial safety and efficacy.^[17] Although eptifibatide has been fairly studied in patients with CAD, there are few available data in the case of a single dose (10 cc) injection prior to coronary stent implantation to justify its use in PCI. The objective of the present study was to evaluate the effect of eptifibatide before intracoronary stent implantation on immediate outcomes in patients with CAD in a double-blind RCT.

MATERIALS AND METHODS

The design of this study was a single-center, double-blind RCT comparing eptifibatide plus stenting with stenting alone in 207 patients during intracoronary stent implantation.

The study was conducted between January and December 2012 and performed in an educational hospital (Shahid-Chamran hospital, Isfahan, Iran). The intervention protocol was approved by the Ethics Committee of the Isfahan University of Medical Sciences, and all patients provided written informed consent. The nonprobability consecutive sampling method was used.

All patients were randomly divided into one of the two groups by Random Allocation Software (SPSS Inc. Chicago).^[18] According to the randomization protocol, eptifibatide (150 µg/kg) was administered in 107 patients just before the stenting procedure.

The inclusion criteria were: Age 30-75 years, stable or unstable angina and/or documentation of myocardial ischemia attributable to coronary artery stenosis, patients without renal failure (renal failure defined as a serum creatinine ≥ 2 mg/dl), and patients without any infectious or systemic disease.

It was decided that cardiac death, stent thrombosis (ST), MI, rates of target lesion and vessel revascularization, cerebral vascular accident (CVA), and emergency CABG would be the major endpoints of the present study.

MI will be distinguished as Q wave or non-Q wave (defined as elevation of total creatine kinase [CK] two times above the upper limit of normal with a positive MB fraction in the absence of pathological Q waves).^[19] Target lesion revascularization (TLR) is defined as any revascularization performed on the treated segment and target vessel revascularization (TVR) as any percutaneous reintervention performed on the treated vessel.^[20] Urgent or emergency CABG is defined as a surgical procedure immediately following catheterization, mandated by clinical circumstances.^[21] Acute coronary ST, defined as ST within 30 days of deployment, has been reduced to <1-2% as a result of improved deployment techniques that fully appose stent to the vessel wall and the use of antiplatelet agents.^[22]

All data were analyzed using PASW 18.0 (SPSS Inc., Chicago, IL, USA). Data are presented as Mean \pm SD for continuous variables and number (percent) for categorical ones.

We used the Kolmogorov-Smirnov test for normality in order to evaluate the assumption of *t*-test. Statistical differences between studied groups were assessed by the independent-samples *t*-test, the Chi-square test, and Fisher's exact test. $P < 0.05$ denoted statistical significance.

RESULTS

The flow-diagram of the study is shown in Figure 1. A total of 110 males (53.1%) and 97 females (46.9%) were included in the analysis. The mean age of the participants was 61.1 ± 8.6 years (range, 45-87 years), and there were no significant statistical differences in age and sex distributions between the two groups ($P = 0.81$ and $P = 0.97$, respectively). The demographic and clinical characteristics of the study population categorized by groups are shown in detail in Table 1.

As is shown in the table, the frequency of smoking, hypertension, family history of CAD, prior MI, and hyperlipidemia were not statistically significant ($P > 0.05$).

The frequency of diabetes mellitus (DM) in the case group was significantly higher than that in the controls (42% vs. 21.5%; $P = 0.001$).

In this study we did not find any significant to statistical differences between the eptifibatide group and the control group in clinical outcomes such as cardiac and noncardiac death, ST, MI, TLR, TVR, CVA, and emergency CABG ($P > 0.05$).

DISCUSSION

Our study examined the clinical immediate outcomes in a consecutive series of patients with CAD who underwent primary PCI and received eptifibatide plus stenting, or stenting alone as a control group, in a double-blind RCT.

The principal finding of the present investigation is that in the studied population of patients with CAD, single-dose intracoronary eptifibatide plus stenting result in similar clinical outcomes when compared to stenting alone.

In accordance with our findings, Raveendran *et al.* in an observational analysis conducted between January 1999 and January 2004 compared the clinical outcomes of 576 patients treated with eptifibatide ($n = 249$) or abciximab ($n = 327$) during primary PCI. As they noted, there were no significant differences between the outcomes of in-hospital death or MI in both eptifibatide- and abciximab-treated groups. They also remarked that eptifibatide was useful and safe as adjunctive pharmacotherapy for patients undergoing primary PCI for acute MI in comparison with abciximab.^[3] In an investigation done between October 2002 and July 2006, different outcomes of 3,541 patients concerning the use of eptifibatide ($n = 2,812$) instead of abciximab ($n = 729$) undergoing primary PCI were assessed. As their results indicated, there were no significant differences in case of early outcomes of patients treated with eptifibatide compared with patients treated with




Figure 1: Flow diagram of the study

Table 1: Demographic features and immediate clinical outcomes of 207 patients divided by studied groups

Characteristics	Groups		P
	Eptifibatide (n=100)	Control (n=107)	
Age (years)	67.2±7.9	66.9±9.2	0.81
Gender (male/female)	47/53	50/57	0.97
History of smoking	32 (32)	28 (26.2)	0.36
DM	24 (42)	23 (21.5)	0.001
HTN	34 (34)	24 (22.4)	0.064
Family history of CAD	23 (23)	14 (13.1)	0.063
Prior MI	11 (11)	10 (9.3)	0.694
HLP	21 (21)	21 (19.6)	0.806
Vessel place			0.018
LAD /LCX /RCA	44/19/37	68/14/25	
Mortality			0.684
Cardiac death	0	2 (1.9)	
Noncardiac death	2 (2)	2 (1.9)	
ST	1 (1)	2 (1.9)	0.601
MI	2 (2)	1 (0.9)	0.522
TLR	1 (1)	2 (1.9)	0.601
TVR	2 (2)	1 (0.9)	0.522
CVA	1 (1)	5 (4.7)	0.115
Emergency CABG	0	0	-

DM: Diabetes mellitus, HTN: Hypertension, CAD: Coronary artery disease, MI: Myocardial infarction, HLP: Hyperlipidemia, LAD: Left anterior descending coronary artery, LCX: Left circumflex coronary artery, RCA: Right coronary artery, ST: Stent thrombosis, TLR: Target lesion revascularization, TVR: Target vessel revascularization, CVA: Cerebral vascular accident, CABG: Coronary artery bypass grafting. Data are expressed as Mean±SD, number and number (percent). P values calculated by independent-sample t-test, chi-square test, and Fisher's exact test

abciximab.^[8] Also, in a study published in 2002 by Stone *et al.*, upon 2,082 patients with acute MI and in a comparison of angioplasty with stenting, with or without abciximab, no significant differences were observed between the percutaneous transluminal coronary angioplasty plus abciximab, stenting alone, or stenting plus abciximab cohorts at 30 days follow-up.^[23] The incorporation of the results given by Stone *et al.* and reports that indicated similar outcomes for eptifibatide and abciximab could be a confirmation of our findings.

The Enhanced Suppression of the Platelet IIb/IIIa Receptor with Integrilin Therapy (ESPRIT) indicated the efficacy of adjunctive eptifibatide therapy during coronary stent implantation at 48 h and at 30 days follow-up. Secondary studies proved the benefits of eptifibatide upon composite rates of death or MI and death, infarction, or TVR during 6 and 12 months follow-up.^[11,15,24]

Although several studies have shown obvious reduction in a variety of ischemic events in patients, which results from receiving eptifibatide as adjunctive pharmacotherapy during PCI,^[25,26] in our results we could not find any significant differences concerning

the effect of eptifibatide upon clinical outcomes compared with the control group. The failure to reach any statistical significance between the two groups (eptifibatide and control), despite the differences reported by former studies, may be related to the small sample size, which greatly reduced the power of the statistical analyses, or to the relatively short time of follow-up. Some supplementary studies with larger sample sizes and more follow-up time are needed to evaluate the real effect of eptifibatide on clinical outcomes.

CONCLUSION

We could not find any significant statistical differences between the short-term clinical outcomes of the single-dose intracoronary eptifibatide and control groups in patients undergoing primary PCI during stent implantation. On the other hand, the use of eptifibatide before intracoronary stent implantation did not have any positive effect on immediate outcomes in patients with CAD.

Acknowledgements

The study was supported by Vice Chancellor for Research (School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran) [Project number: 391460].

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Skoog M, Grevstad B, Wetterslev J, Hróbjartsson A, Gluud C. Antibiotics for secondary prevention of coronary heart disease. The Cochrane Library 2013.
2. Htay T, Liu MW. Drug-eluting stent: A review and update. *Vasc Health Risk Manag* 2005;1:263-76.
3. Raveendran G, Ting HH, Best PJ, Holmes DR Jr, Lennon RJ, Singh M, *et al.* Eptifibatide vs abciximab as adjunctive therapy during primary percutaneous coronary intervention for acute myocardial infarction. *Mayo Clin Proc* 2007;82:196-202.
4. Boersma E, Harrington RA, Moliterno DJ, White H, Théroux P, Van de Werf F, *et al.* Platelet glycoprotein IIb/IIIa inhibitors in acute coronary syndromes: A meta-analysis of all major randomised clinical trials. *Lancet* 2002;359:189-98.
5. Anderson KM, Califf RM, Stone GW, Neumann FJ, Montalescot G, Miller DP, *et al.* Long-term mortality benefit with abciximab in patients undergoing percutaneous coronary intervention. *J Am Coll Cardiol* 2001;37:2059-65.
6. Randomised placebo-controlled trial of abciximab before and during coronary intervention in refractory unstable angina: The CAPTURE Study. *Lancet* 1997;349:1429-35.
7. Vorchheimer DA, Badimon JJ, Fuster V. Platelet glycoprotein IIb/IIIa receptor antagonists in cardiovascular disease. *JAMA* 1999;281:1407-14.
8. Gurm HS, Smith DE, Collins JS, Share D, Riba A, Carter AJ, *et al.*; Blue Cross Blue Shield of Michigan Cardiovascular Consortium (BMC2).

- The relative safety and efficacy of abciximab and eptifibatide in patients undergoing primary percutaneous coronary intervention: Insights from a large regional registry of contemporary percutaneous coronary intervention. *J Am Coll Cardiol* 2008;51:529-35.
9. Horne BD, Knight S, May HT. Panoptic total cardiovascular risk prediction using all predictors versus optimized risk assessment using variable subsets. *Future Cardiol* 2012;8:765-78.
 10. Kleiman NS, Lincoff AM, Flaker GC, Pieper KS, Wilcox RG, Berdan LG, *et al.* Early percutaneous coronary intervention, platelet inhibition with eptifibatide, and clinical outcomes in patients with acute coronary syndromes. PURSUIT Investigators. *Circulation* 2000;101:751-7.
 11. ESPRIT Investigators. Enhanced Suppression of the Platelet IIb/IIIa Receptor with Integrilin Therapy. Novel dosing regimen of eptifibatide in planned coronary stent implantation (ESPRIT): A randomised, placebo-controlled trial. *Lancet* 2000;356:2037-44.
 12. Zeymer U, Zahn R, Schiele R, Jansen W, Girth E, Gitt A, *et al.* Early eptifibatide improves TIMI 3 patency before primary percutaneous coronary intervention for acute ST elevation myocardial infarction: Results of the randomized integrilin in acute myocardial infarction (INTAMI) pilot trial. *Eur Heart J* 2005;26:1971-7.
 13. Dery JP, Campbell ME, Mathias J, Pieper KS, Harrington RA, Madan M, *et al.* ESPRIT Investigators. Complementary effects of thienopyridine pretreatment and platelet glycoprotein IIb/IIIa integrin blockade with eptifibatide in coronary stent intervention; results from the ESPRIT trial. *Catheter Cardiovasc Interv* 2007;70:43-50.
 14. Marmur JD, Poludasu S, Lazar J, Cavusoglu E. Long-term mortality after bolus-only administration of abciximab, eptifibatide, or tirofiban during percutaneous coronary intervention. *Catheter Cardiovasc Interv* 2009;73:214-21.
 15. O'Shea JC, Hafley GE, Greenberg S, Hasselblad V, Lorenz TJ, Kitt MM, *et al.* ESPRIT Investigators (Enhanced Suppression of the Platelet IIb/IIIa Receptor with Integrilin Therapy trial) Platelet glycoprotein IIb/IIIa integrin blockade with eptifibatide in coronary stent intervention: The ESPRIT trial: A randomized controlled trial. *JAMA* 2001;285:2468-73.
 16. Deibele AJ, Jennings LK, Tcheng JE, Neva C, Earhart AD, Gibson CM. Intracoronary eptifibatide bolus administration during percutaneous coronary revascularization for acute coronary syndromes with evaluation of platelet glycoprotein IIb/IIIa receptor occupancy and platelet function: The Intracoronary Eptifibatide (ICE) Trial. *Circulation* 2010;121:784-91.
 17. Midei MG, Coombs VJ, Lowry DR, Drossner MN, Prewitt KC, Wang JC, *et al.* Clinical outcomes comparing eptifibatide and abciximab in ST elevation acute myocardial infarction patients undergoing percutaneous coronary interventions. *Cardiology* 2006;107:172-7.
 18. Saghaei M. Random allocation software for parallel group randomized trials. *BMC Med Res Methodol* 2004;4:26.
 19. Biondi-Zoccai GG, Valgimigli M, Sheiban I, Margheri M, Marzocchi A, Prati F, *et al.* A randomized trial comparing eptifibatide vs. placebo in patients with diffuse coronary artery disease undergoing drug-eluting stent implantation: Design of the INtegrilin plus STenting to Avoid myocardial Necrosis Trial. *J Cardiovasc Med (Hagerstown)* 2008;9:957-62.
 20. Biondi-Zoccai GG, Sangiorgi GM, Antoniucci D, Grube E, Di Mario C, Reimers B, *et al.* Taxus in Real-life Usage Evaluation Study. Testing prospectively the effectiveness and safety of paclitaxel-eluting stents in over 1000 very high-risk patients: Design, baseline characteristics, procedural data and in-hospital outcomes of the multicenter Taxus in real-life usage evaluation (TRUE) study. *Int J Cardiol* 2007;117:349-54.
 21. Aldea GS. Use of heparin-bonded cardiopulmonary bypass circuits with alternatives to standard anticoagulation. *Advanced Therapy in Cardiac Surgery* 2003;103:46.
 22. Farb A, Burke AP, Kolodgie FD, Virmani R. Pathological mechanisms of fatal late coronary stent thrombosis in humans. *Circulation* 2003;108:1701-6.
 23. Stone GW, Grines CL, Cox DA, Garcia E, Tcheng JE, Griffin JJ, *et al.* Controlled Abciximab and Device Investigation to Lower Late Angioplasty Complications (CADILLAC) Investigators. Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N Engl J Med* 2002;346:957-66.
 24. O'Shea JC, Buller CE, Cantor WJ, Chandler AB, Cohen EA, Cohen DJ, *et al.* ESPRIT Investigators. Long-term efficacy of platelet glycoprotein IIb/IIIa integrin blockade with eptifibatide in coronary stent intervention. *JAMA* 2002;287:618-21.
 25. Granada JF, Kleiman NS. Therapeutic use of intravenous eptifibatide in patients undergoing percutaneous coronary intervention: Acute coronary syndromes and elective stenting. *Am J Cardiovasc Drugs* 2004;4:31-41.
 26. O'Shea JC, Tcheng JE. Eptifibatide: A potent inhibitor of the platelet receptor integrin glycoprotein IIb/IIIa. *Expert Opin Pharmacother* 2002;3:1199-210.