

Case Report

Isolated pulmonary candidiasis in a patient with diabetes mellitus: A rare case report

Atousa Hakamifard, Farzin Khorvash¹, Arash Raisi²

Infectious Diseases and Tropical Medicine Research Center, Isfahan University of Medical Sciences, ¹Nosocomial Infection Research Center, Isfahan University of Medical Sciences, ²Department of Hematology, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract Fungal infections are as a cause of morbidity and mortality in immunocompromise patients. Because the respiratory tract is colonized with *Candida*, the presence of this agent in respiratory specimens makes the diagnosis of *Candida* pneumonia problematic. *Candida* pneumonia is a rare infection, and the majority of cases are secondary to hematogenous dissemination. Furthermore isolated *Candida* pneumonia originating from endotracheal inoculation is an extremely rare entity. We describe a case of isolated pulmonary candidiasis in the form multiple nodular lesions in a patient with long-term history of diabetes mellitus without evidence of fungemia or systemic involvement who responded to antifungal therapy.

Key Words: *Candida*, Candidiasis, fluconazole, pneumonia

Address for correspondence:

Prof. Farzin Khorvash, Nosocomial Infection Research Center, Isfahan University of Medical Sciences, Isfahan, Iran. E-mail: khorvash@med.mui.ac.ir

Received: 27.10.2015, **Accepted:** 26.12.2015

INTRODUCTION

A genus of yeasts as the significant cause of fungal infections worldwide is *Candida*.^[1] The most prevalent *Candida* species is *Candida albicans*. This fungus is a part of the normal flora of upper respiratory and gastrointestinal tract.^[2] However, *Candida* pneumonia is a rare manifestation, especially primary form.^[3,4] Most of the cases of *Candida* pneumonia occurs in immunosuppressed patients with hematogenous dissemination from a distant site.^[4] There are several risk factors for candidiasis such as the broad spectrum of antibiotics, neutropenia, indwelling catheter, diabetes mellitus (DM), human immunodeficiency virus (HIV), acute renal failure, surgical procedure, and long term corticosteroid treatment. Predisposing factors for pulmonary candidiasis also are the same.

Fungal colonization is considered a major risk factor for subsequent fungal disease in these patients.^[4,5]

In this paper, we report a rare case of isolated *Candida* pneumonia in the form of nodular lesions in a patient with DM that responded to antifungal therapy.

CASE REPORT

A 55-year-old man with a 10-year history of DM was admitted to our hospital with a 2 months history of dyspnea and cough. The patient was alert and afebrile and complained of chronic nonproductive cough. The patient had no history of immunosuppressive treatment or disease except the history of DM. The patient denied alcohol abuse and smoking. His vital sign was stable. Physical examination was not notable. Cell blood count (CBC) showed hemoglobin (Hgb) 9.9g/dl, white blood cell count (WBC)

Access this article online	
Quick Response Code:	Website: www.advbiores.net
	DOI: 10.4103/2277-9175.187377

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Hakamifard A, Khorvash F, Raisi A. Isolated pulmonary candidiasis in a patient with diabetes mellitus: A rare case report. *Adv Biomed Res* 2016;5:146.

6500, with polymorphonuclear dominant (56%). The level of C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR) were 50mg/dl and 105 mm/hour respectively. HIV test was negative. In chest X-ray, nodular pattern was seen. This pattern also was showed in thorax computed tomography (CT) scan [Figure 1].

Considering the course of the disease, clinical manifestations and stable vital signs, the antibiotic was not started.

3 sets of blood cultures and echocardiogram were negative for endocarditis. Examination of the sputum specimen showed *C.albicans*. As the patient complained of a chronic cough and according to pulmonary lesions, bronchoscopy and bronchoalveolar lavage (BAL) was performed. The specimens evaluated for bacteria, nocardia, actinomyces, fungi and mycobacterium tuberculosis (TB) smear and culture, polymerase chain reaction for TB and also cytology for malignancies. The results were all negative except the culture of BAL fluid which revealed *C. albicans*. The serum and BAL fluid galactomannan were negative.

Hence, CT-guided percutaneous biopsy of the lesion of the lung was performed. The result showed yeast and pseudohyphae that was compatible with the diagnosis of *Candida*. The tissue culture yielded *C. albicans* as well. Furthermore, microscopic studies for other microorganism such as acid-fast staining were negative. On special sectioning, no evidence of granuloma or malignancy was reported. Considering diagnosis, antifungal regimen (intravenous fluconazole) was started. Parenteral antifungal therapy continued for 2 weeks and then followed with oral fluconazole. Control chest CT scan was done 5 weeks later, which showed significant regression of pulmonary

lesions [Figure 2]. At the completion of treatment, the level of CRP and ESR decrease to 6 mg/dl and 17mm/hr, respectively.

DISCUSSION

In this paper, we describe a patient with isolated *Candida* pneumonia. Our patient represents *Candida* pneumonia with an unusual manifestation. Pulmonary candidiasis is a rare entity often in immune deficient patients with the majority of cases secondary to fungemia. Because the respiratory tract is colonized with *Candida*, the presence of this agent in respiratory specimens makes the diagnosis of pneumonia problematic which represents contamination versus disease.^[4] Although BAL culture is an important adjunctive method,^[6] (as in our case with positive BAL culture) but the growth of *Candida* in respiratory samples represents contamination; hence, the diagnosis depends mainly on tissue biopsy with evidence of fungal invasion.^[4] Predisposing factors are immunodeficiency states such as immunosuppressive diseases, malignancies and corticosteroids.^[7] The patient with a history of DM was described in this paper. DM has been introduced as a predisposing factor for colonization and infection.^[8,9] Hence, this might have been responsible for the increased risk of infection. In this case, multiple nodular lesions were seen and the tissue biopsy confirmed the diagnosis of *Candida*. *Candida* pneumonia originating from endobronchial inoculation of the lung is extremely rare and usually is in the form of diffuse or lobar bronchopneumonia. On the contrary, hematogenous dissemination results in multiple nodular lesions of the lung.^[4] One of the interesting aspects of this case was the isolated lung infection in the form of multiple pulmonary nodules without evidence of fungemia or other systemic infections which are rare and describe in a few case




Figure 1: Multiple nodular lesions with ground glass halo



Figure 2: All of the lesions significantly were resolved after treatment

reports.^[6,3] In another study by Yasuda *et al.*, *Candida* pneumonia with multiple cavitory lesions and nodules was described as a case of invasive candidiasis, and the blood culture specimens yielded *C.albicans*.^[10] In our case intravenous fluconazole resulted in clinical improvement and also diminished inflammatory markers.

In conclusion, although candida pneumonia is rare, it should be considered in differential diagnosis of pneumonia in the proper setting and for definitive diagnosis the tissue biopsy should be obtained.

Acknowledgment

We gratefully acknowledge all the academic staff of Infectious Diseases Department and Hematology Department of Isfahan University of Medical Sciences.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Manolakaki D, Velmahos G, Kourkoumpetis T, Chang Y, Alam HB, De Moya MM, *et al.* *Candida* infection and colonization among trauma patients. *Virulence* 2010;1:367-75.
2. Nobile CJ, Johnson AD. *Candida albicans* biofilms and human disease. *Annu Rev Microbiol* 2015;69:71-92.
3. Shweihat Y, Perry J III, Shah D. Isolated *Candida* infection of the lung. *Respir Med Case Rep* 2015;16:18-19.
4. Edwards JE Jr. *Candida* species. Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases. 8th ed. Philadelphia: Churchill Livingstone; 2015. p. 2879-94.
5. Casqueiro J, Casqueiro J, Alves C. Infections in patients with diabetes mellitus: A review of pathogenesis. *Indian J Endocrinol Metab* 2012; 16 Suppl 1:S27-36.
6. Knox KS, Meinke L. Role of bronchoalveolar lavage diagnostics in fungal infections. *Clin Chest Med* 2009;30:355-65.
7. Yapar N. Epidemiology and risk factors for invasive candidiasis. *Ther Clin Risk Manag* 2014;10:95-105.
8. Leroy O, Gangneux JP, Montravers P, Mira JP, Gouin F, Sollet JP, *et al.* Epidemiology, management, and risk factors for death of invasive *Candida* infections in critical care: A multicenter, prospective, observational study in France (2005-2006). *Crit Care Med* 2009;37:1612-8.
9. Aly FZ, Blackwell CC, Mackenzie DA, Weir DM, Clarke BF. Factors influencing oral carriage of yeasts among individuals with diabetes mellitus. *Epidemiol Infect* 1992;109:507-18.
10. Yasuda Y, Tobino K, Asaji M, Yamaji Y, Tsuruno K. Invasive candidiasis presenting multiple pulmonary cavitory lesions on chest computed tomography. *Multidiscip Respir Med* 2015;10:11.