

Cytotoxic, antioxidant and phytochemical analysis of *Gracilaria* species from Persian Gulf

Alireza Ghannadi, Leila Shabani¹, Afsaneh Yegdaneh

Department of Pharmacognosy, School of Pharmacy and Pharmaceutical Sciences and Isfahan Pharmaceutical Sciences Research Center, Isfahan University of Medical Sciences, Isfahan, ¹Department of Biology, Faculty of Sciences, Shahrekord University, Shahrekord, Iran

Abstract

Background: Marine algae, also called seaweeds, are abundantly present in the coastal area of Iran, especially in Persian Gulf. These plants contain important phytochemical constituents and have potential biological activities. The present study investigated the presence of phytochemical constituents and total phenolic quantification of the seaweeds *Gracilaria salicornia* and *Gracilaria corticata*. Cytotoxicity of seaweeds was tested against HT-29, HeLa, and MCF-7 cell lines. Antioxidant potential of these two *Gracilaria* species was also analyzed.

Materials and Methods: Extracts of *G. salicornia* and *G. corticata* were subjected to phytochemical and cytotoxicity tests. Phytochemical screenings were employed to identify the chemical constituents and total phenolic content. Cytotoxicity was characterized by IC₅₀ of human cancer cell lines (MCF-7, HeLa, and HT-29) using sulforhodamine assay. Antioxidant activities were evaluated using 2,2-diphenyl-1-picrylhydrazyl.

Results: The analysis revealed that tannins were the most abundant compounds in *G. corticata* while sterols and triterpenes were the most abundant ones in *G. salicornia*, but the total phenolic content of the two seaweeds was similar. Cytotoxic results showed that both species could inhibit cell growth effectively, especially against HT-29 cell line.

Conclusion: Considerable phytochemicals, high antioxidant potential, and moderate cytotoxic activity of *G. salicornia* and *G. corticata* make them appropriate candidates for further studies and identification of their bioactive principles.

Key Words: Antioxidant, cytotoxic, *Gracilaria corticata*, *Gracilaria salicornia*, Persian Gulf, phytochemical, seaweed

Address for correspondence:

Dr. Afsaneh Yegdaneh, Department of Pharmacognosy, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran. E-mail: yekdaneh@pharm.mui.ac.ir

Received: 14.07.2015, **Accepted:** 17.11.2015

INTRODUCTION

The unique field of marine natural products chemistry emerged in the late 1960s, developed rapidly during the 1980s, and matured in the last decade. The marine biosphere is a rich source of biologically-active

principles originating from marine organisms such as seaweeds, invertebrates, coral reefs and marine bacteria. As a result of the potential for new drug discovery, marine natural products have attracted scientists from different disciplines. This interest has led to the discovery of thousands of secondary

Access this article online	
Quick Response Code:	Website: www.advbiores.net
	DOI: 10.4103/2277-9175.187373

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Ghannadi A, Shabani L, Yegdaneh A. Cytotoxic, antioxidant and phytochemical analysis of *Gracilaria* species from Persian Gulf. *Adv Biomed Res* 2016;5:139.

metabolites from marine organisms, many of which have shown potent biological activities.^[1-3]

Seaweeds belong to a group of plants known as algae. Seaweeds are classified as Rhodophyta (red algae), Phaeophyta (brown algae), and Chlorophyta (green algae) depending on their nutrient and chemical composition. Like other plants, seaweeds contain various inorganic and organic substances which can benefit human health.^[4] Algae contain amino acids, terpenoids, phlorotannins, steroids, phenolic compounds, halogenated ketones and alkanes and cyclic polysulphides.^[5] They are rich sources of structurally novel and biologically active metabolites. The metabolites, both primary and secondary, produced by seaweeds may be potential bioactive compounds, which are used in pharmaceutical industry.^[6] Marine algae have been used as medicine for a long time^[7] and have been extensively studied by several researchers.^[2,3,8] They are the sustainable resources in the marine ecosystem and are used as a source of food, fodder, fertilizer, industrial productions, treatment of effluents^[9] and medicine. Recent studies revealed that seaweeds contain antibacterial,^[10] antiviral,^[11,12] antifungal,^[13] cytotoxic^[14] and acetylcholinesterase inhibitor^[15] properties.

Iran has coastal lines about 1260 km along the Persian Gulf and the Oman Sea. More than 250 species of different alga have been identified in this area among which from Chlorophyta: Ulvaceae and Caulerpaceae;^[16,17] from Phaeophyta: Dictyotaceae and Sargassaceae; and from Rhodophyta: Gracilariaceae, Gelidiaceae, and Hypneaceae are the most abundant families.^[18] Despite the large extent of marine algae in this region, there are only a few studies about phytochemical and biological analysis of these seaweeds. The present study aims to identify types of active constituents of the seaweeds *Gracilaria corticata* and *Gracilaria salicornia*, antioxidant potential, and also cytotoxic activity of extracts against MCF-7, HeLa, and HT-29 cell lines.

MATERIALS AND METHODS

Authentication of plant material

The seaweeds were collected from the Persian Gulf coasts of Iran, Bushehr Province, in autumn 2014. Voucher specimens were made and deposited in the herbarium of the School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences (Code: 2666, 2667) and were identified by Agricultural and Natural Resources Research Center of Bushehr.

Preparation of the extracts

The plant samples were cut into small pieces and completely air-dried, and stored in glass containers until extraction. About 100 g of the dried plant material was extracted for 5 days with methanol. The extracts were filtered through two layers of cotton fabric and evaporated at room temperature, under reduced pressure, to a dry residue and stored in sterile vial pending phytochemical and cytotoxic screening.

In vitro cytotoxicity assay

The extracts were tested on MCF-7 (human breast adenocarcinoma), HeLa (cervical carcinoma), and HT-29 (human colon adenocarcinoma) cells. The cancer cell lines were purchased from the National Cell Bank of Pasteur Institute of Iran and grown in Dulbecco's Modified Eagle Medium (D-MEM) supplemented with 10% fetal bovine serum. Cells were seeded in 96-wells (3500 cells/well) and allowed to adhere for 24 h at 37°C, with 5% CO₂ in a fully humidified incubator. Then, 100 µl of serially diluted samples in medium were dispensed into the wells of the cell plates and incubated for a further 72 h. After removal of the sample medium, the cells were topped up with 200 µl D-MEM medium and incubated. After 72 h cells were fixed with cold 40% trichloroacetic acid at 4°C for 1 h and washed with tap water. These cells were determined by sulforhodamine assay.^[19] The absorbance was measured at 492 nm using a microplate reader. Percentage of dead cells was calculated in comparison to control. The concentration of the extract that inhibited 50% cells growth (IC₅₀) was determined from the graph plotted by the concentration versus percentage of dead cells.^[19]

Phytochemical screening

Tests for phytochemical constituents – alkaloids, steroids and triterpenes, anthraquinones, flavonoids, saponins, cyanogenic glycosides, cardiac glycosides and tannins-followed the methods described previously.^[20]

Determination of total phenolics

The powdered plant material (20 g of each sample) was weighed into a 50 ml flask, extracted with 30 ml of ethanol 40% using the sonicator for about 30 min and shaking for about 10 min. After allowing the extracts to cool down to room temperature, the flasks were filled to full volume with extraction solvent.

Preparation of standard

Twenty milligrams of gallic acid and 30 ml 40% EtOH were added into a 50 ml volumetric flask and sonicated until no solid was present in the flask. After allowing the solution to cool down to room temperature, the flask was filled with extraction solvent. The standard solution was diluted several times.

One milliliter of standard solution was transferred to 100 ml volumetric flask with 60–70 high-performance liquid chromatography (HPLC) grade water. The contents swirled to mix. Five milliliter of Folin-Ciocalteu's phenol reagent was added and mixed again. After 1 min and before 8 min, 15 ml of sodium carbonate solution was added, the time recorded as time zero. The volume was made up to 100 ml exactly with HPLC grade water. The flask stoppered and mixed thoroughly by inverting it several times. After 2 h, the ultraviolet absorption range at 550–850 nm and maximum absorbance about 760 nm were recorded. The same solution without the extraction solution was used as blank solution.^[21]

2,2-diphenyl-1-picrylhydrazyl free radical scavenging assay

The free radical scavenging activity was measured using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay. Sample stock solutions (1.0 mg/ml) of the extracts were diluted to final concentrations of 243, 81, 27, 9, 3, and 1 µg/ml in ethanol. One ml of a 50 µg/ml DPPH ethanol solution was added to 2.5 ml of sample solutions of different concentrations and allowed to react at room temperature. After 30 min, the absorbance values were measured at 518 nm and converted into the percentage antioxidant activity (AA) using the following formula:

$$AA\% = \frac{[\text{absorbance of the control} - \text{absorbance of sample}]}{\text{absorbance of the control}} \times 100$$

Ethanol (1.0 ml) plus plant extracts solutions (2.5 ml) were used as a blank solution.

DPPH solution (1.0 ml) plus ethanol (2.5 ml) was used as negative control. The positive controls (ascorbic acid, butylated hydroxyanisole, and butylated hydroxytoluene) were used as standard solutions. Assays were carried out in triplicate.^[22]

RESULTS AND DISCUSSION

Phytochemical constituents

Phytochemical data show distinct patterns of chemical compositions in constituents of the extracts. The patterns of composition differed considerably in their quantitative values. The results of phytochemical evaluation are shown in Table 1.

This analysis showed the most abundant compounds in *G. corticata* were tannins, saponins, sterols, and triterpenes followed by alkaloids, flavonoids, and cardiac glycosides. Cyanogenic glycosides were absent in this seaweed. In *G. corticata*, tannins

were the most abundant compounds followed by saponins, alkaloids, sterols, and triterpenes. Cyanogenic glycosides and anthraquinones were not present.

The amount of total phenol was determined with the Folin–Ciocalteu reagent. Gallic acid was used as a standard compound, and the total phenols were expressed as mg/g gallic acid equivalent using the standard curve equation:

$$y = 1.771x - 0.0252, R^2 = 0.9958.$$

Where y is absorbance at 760 nm and x is total phenolic content in the extracts of different alga expressed in mg/L [Figure 1]. Phenolic compounds are a class of antioxidant agents which act as free radical terminators.^[13] Table 2 shows the contents of total phenols that were measured by Folin–Ciocalteu reagent in terms of gallic acid equivalent. The total phenol in selected seaweeds was 0.012–0.013 mg in 1 g dry extract in *G. salicornia* and *G. corticata*, respectively.

Table 1: Phytochemical constituents of *Gracilaria corticata* and *Gracilaria salicornia*

Constituents	<i>G. salicornia</i> extract	<i>G. corticata</i> extract	Method
Tannins	+++	+++	FeCl ₃ test
Alkaloids	++	++	Hager's test
Dragendorff's test			
Saponins	+++	++	Foam test
Sterols and triterpenes	+++	++	Liebermann–Burchard test
Flavonoids	++	+	Alkaline reagent test
Cyanogenic glycosides	-	-	Wilstatter
Cardiac glycosides	+	+	Keller Kelliani's test
Anthraquinones	+	-	Borntrager's test

G. corticata: *Gracilaria corticata*, *G. salicornia*: *Gracilaria salicornia*




Figure 1: Standard curve of gallic acid

Cytotoxic assay

The criteria used to categorize the activity of extracts against human breast, cervix, colon (MCF-7, HeLa, and HT-29) cell lines based on IC_{50} values were modified from those of NCI and Geran *et al.*^[20] as follows: $IC_{50} \leq 20 \mu\text{g/ml}$ = highly active, $IC_{50} 21\text{--}200 \mu\text{g/ml}$ = moderately active, $IC_{50} 201\text{--}500 \mu\text{g/ml}$ = weakly active, and $IC_{50} > 501 \mu\text{g/ml}$ = inactive.

The cytotoxicity data for the extracts against MCF-7 (human breast adenocarcinoma), HeLa (cervical carcinoma), and HT-29 (human colon adenocarcinoma) cells are displayed in Table 3.

Antioxidant activity

The AA of the seaweed extracts was measured on the basis of the scavenging activity of the stable DPPH free radical. IC_{50} value is inversely related to AA. The extracts of the two algae were found to have different levels of AA. Table 4 shows that higher AA was found in *G. salicornia* extract.

So far more than 2400 marine natural products have been isolated from seaweeds of subtropical and tropical populations.^[23] Certain algae have long been used in traditional Chinese herbal medicine in the treatment of cancer.^[24] Many studies have been developed to determine the bioactive compounds produced by marine algae. Antioxidant and cytotoxic activities are one of the most important specificities of marine algae. Some metabolites such as bromophenols, carotene, and steroids were isolated and purified from some algae, and their activity against some cancer cell lines were demonstrated.^[25]

Table 2: Total phenolic content of seaweeds

Sample	$\lambda_{\text{max}}=760 \text{ nm}$	Concentration (mg/g)
<i>G. salicornia</i>	0.123	0.012
<i>G. corticata</i>	0.136	0.013

G. corticata: *Gracilaria corticata*, *G. salicornia*: *Gracilaria salicornia*

Table 3: Cytotoxic activity (IC_{50}) of the seaweed *Gracilaria corticata* and *Gracilaria salicornia*

Sample	IC_{50} ($\mu\text{g/ml}$)		
	HT-29	HeLa	MCF-7
<i>G. salicornia</i>	68.2	125.9	185.8
<i>G. corticata</i>	58.6	117.4	120.6

G. corticata: *Gracilaria corticata*, *G. salicornia*: *Gracilaria salicornia*

Table 4: Antioxidant activity (IC_{50}) of the seaweed *Gracilaria corticata* and *Gracilaria salicornia*

Sample	IC_{50} (mg/ml)
<i>G. salicornia</i>	0.73
<i>G. corticata</i>	0.54

G. corticata: *Gracilaria corticata*, *G. salicornia*: *Gracilaria salicornia*

In the case of red seaweeds, five major edible genera have been introduced, including *Porphyra*, *Palmaria*, *Gracilaria*, *Gelidium*, and *Kappaphycus* (*Euचेuma*). In particular, numerous red seaweeds of the genus *Gracilaria* are utilized as fresh food in many parts of the world.^[26,27] In this research, an effort was made to study the similarity and differences between the two species on the basis of phytochemical analysis and bioevaluation. Phytochemical tests revealed lots of similarities than differences. The only difference between the two plants was about the detection of anthraquinones which were not present in *G. corticata*. The amount of tannins, sterols, triterpenes, saponins, and flavonoids were also different, but the phenolic content of these seaweeds were similar. Regarding cytotoxic activity, both species showed better results against HT-29 cell lines as compared to MCF-7 and HeLa cell lines, but *G. salicornia* had better results. In a study about cytotoxic activities often different algae from the Persian Gulf and the Oman Sea, *G. salicornia* was tested against MDA-MB-231, MCF-7, and T-47D cell lines but IC_{50} of this alga was reported higher than $400 \mu\text{g/ml}$.^[28] The difference between our data and this study may be because of different solvents used for extraction, season and time of collection, and also different methods. Saeidnia *et al.* also reported that ethyl acetate extract of *G. salicornia* showed a potent cytotoxic effect against *Artemia salina nauplii*.^[29] Phytochemical analysis showed that sterols and triterpenes are the main compounds in the ethyl acetate fraction and may be responsible for the cytotoxic activity of this seaweed. Antioxidant capacity is widely used as a parameter for medicinal bioactive components. Natural antioxidants are known to exhibit a wide range of biological effects including antibacterial, antiviral, anti-inflammatory, antiallergic, antithrombotic, and vasodilatory activities.^[30] The present study infers that *G. salicornia* and *G. corticata* extracts exhibit effective antitumor activity and seem to have potent secondary metabolites. They are cost-effective, easy to produce and purify. In future, they can be recommended to patients as an effective therapeutic tool in the form of a food or drug. Further research needs to be explored to study the bioactive compounds of these two seaweeds and for the successful implication of them as a potent therapeutic tool against cancer.

CONCLUSION

The extracts of two seaweeds from Persian Gulf, Iran were screened for their antioxidant, cytotoxic and phytochemical analysis. Phytochemical and cytotoxic results were similar with a few differences about all species. These seaweed extracts and their active components could emerge as natural and

alternative antioxidants or serve as starting points for synthesizing more effective cytotoxic drugs.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Smit AJ. Medicinal and pharmaceutical uses of seaweed natural products: A review. *J Appl Phycol* 2004;16:245-62.
2. Yegdaneh A, Putschakarn S, Yuenyongsawad S, Ghannadi A, Plubrukarn A. 3-oxoabolene and 1-oxocurcuphenol, aromatic bisabolanes from the sponge *Myrmekioderma* sp. *Nat Prod Commun* 2013;8:1355-7.
3. Blunt JW, Copp BR, Hu WP, Munro MH, Northcote PT, Prinsep MR. Marine natural products. *Nat Prod Rep* 2007;24:31-86.
4. Kuda T, Taniguchi E, Nishizawa M, Araki Y. Fate of water-soluble polysaccharides in dried *Chorda filum* a brown alga during water washing. *J Food Compost Anal* 2002;15:3-9.
5. Mtolera MS, Semesi AK. Antibacterial activity of extracts from six green algae from Tanzania. In: *Current Trends in Marine Botanical Research in the East African Region*. Uppsala, Sweden: Gotab AB; 2000. p. 211-7.
6. Kusumoto IT, Nakabayashi T, Kida H, Miyashiro T, Hattori M, Namba T. Shimotohno Screening of various plant extracts used in ayurvedic medicine for inhibitory effects on human immunodeficiency virus-I (HIV-I) protease. *Phytother Res* 1995;9:180.
7. Fitton JH. Antiviral properties of marine algae. In: Critchley AT, Ohno M, Largo DB, editors. *World Seaweed Resources*. Wokingham, UK: Windows and Macintosh, ETI Information Services; 2006. p. 7.
8. González del Val A, Platas G, Basilio A, Cabello A, Gorrochategui J, Suay I, *et al.* Screening of antimicrobial activities in red, green and brown macroalgae from Gran Canaria (Canary Islands, Spain). *Int Microbiol* 2001;4:35-40.
9. Sharmila S, Rebecca LJ. A comparative study on the degradation of leather industry effluent by marine algae. *Int J Pharm Sci Rev Res* 2014;25:46-50.
10. Tuney I, Cadirci BH, Unal D, Sukatar A. Antimicrobial activities of the extracts of marine algae from the coast of Urla (Izmir, Turkey). *Turk J Biol* 2006;30:171-5.
11. Garg HS, Sharma T, Bhakuni DS, Pramanik BN, Bose AK. An antiviral sphingosine derivative from green alga *Ulva fasciata*. *Tetrahedron Lett* 1992;33:1641-4.
12. Tang HF, Yang-Hua Y, Yao XS, Xu QZ, Zhang SY, Lin HW. Bioactive steroids from the brown alga *Sargassum carpophyllum*. *J Asian Nat Prod Res* 2002;4:95-101.
13. Aliya R, Shameel M. Phycochemical evaluation of four coenocytic green seaweeds from the coast of Karachi. *Pak J Mar Biol* 1999;5:65-76.
14. Thangam TS, Kathiresan K. Mosquito larvicidal activity of marine plant extracts with synthetic insecticides. *Bot Mar* 1991;34:537-9.
15. Ghannadi A, Plubrukarn A, Zandi K, Sartavi K, Yegdaneh A. Screening for antimalarial and acetylcholinesterase inhibitory activities of some Iranian seaweeds. *Res Pharm Sci* 2013;8:113-8.
16. Sohrabipour J, Rabiei R. The checklist of green algae of the Iranian coastal lines of the Persian Gulf and Gulf of Oman. *Iran J Bot* 2007;13:146-9.
17. Sohrabipour J, Nejadsatari T, Assadi M, Rebei R. The marine algae of the Southern coast of Iran, Persian Gulf, Lengeh area. *Iran J Bot* 2004;10:83-93.
18. Safa O, Soltanipoor MA, Rastegar S, Kazemi M, Nourbakhsh Dehkordi K, Ghannadi A. An ethnobotanical survey on hormozgan province, Iran. *Avicenna J Phytomed* 2013;3:64-81.
19. Kamba AS, Hassan LG. Phytochemical screening and antimicrobial activities of *Euphorbia balsamifera* leaves, stems and root against some pathogenic microorganisms. *Afr J Pharm Pharmacol* 2010;4:645-52.
20. Geran RI, Greenberg NH, Macdonald MM, Shumacher AM, Abbott BJ. Protocols for screening chemical agents and natural products against animal tumors and other biological systems. *Cancer Chem Rep* 1972;3:1-103.
21. Pourmorad F, Hosseinimehr SJ, Shahabimajid N. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr J Biotechnol* 2006;5:1142-5.
22. Braca A, De Tommasi N, Di Bari L, Pizzi C, Politi M, Morelli I. Antioxidant principles from *Bauhinia tarapotensis*. *J Nat Prod* 2001;64:892-5.
23. Manilal A, Sujith S, Kiran GS, Selvin J, Shakir C, Gandhimathi R, *et al.* Biopotentials of seaweeds collected from South West Coast of India. *J Mar Sci Technol* 2009;17:67-73.
24. Yamamoto I, Takahashi M, Tamura E, Maruyama H, Mori H. Antitumor activity of edible marine algae: Effect of crude fucoidan fractions prepared from edible brown seaweeds against L-1210 leukemia. *Hydrobiologia* 1984;116:145-8.
25. Xu N, Fan X, Yan X, Tseng C. Screening marine algae from China for their antitumor activities. *J Appl Phycol* 2004;16:451-6.
26. McDermid KJ, Stuercke B. Nutritional composition of edible Hawaiian seaweeds. *J Appl Phycol* 2003;15:513-24.
27. Norziah MH, Ching CY. Nutritional composition of edible seaweed *Gracilaria changgi*. *Food Chem* 2006;68:69-76.
28. Erfani N, Nazemosadat Z, Moein M. Cytotoxic activity of ten algae from the Persian Gulf and Oman Sea on human breast cancer cell lines; MDA-MB-231, MCF-7, and T-47D. *Pharmacognosy Res* 2015;7:133-7.
29. Saeidnia S, Gohari AR, Shahverdi AR, Permeh P, Nasiri M, Mollazadeh K. *In vitro* antitumor activity of *Gracilaria corticata* (a red alga) against jurkat and molt-4 human cancer cell lines. *Afr J Biotechnol* 2010;9:6787-90.
30. Cook NC, Samman S. Flavonoids: Chemistry, metabolism, cardioprotective effects and dietary sources. *J Nutr Biochem* 1996;7:66-76.