

# Chromium level in prediction of diabetes in pre-diabetic patients

Rahmatollah Rafiei, Zahra Habyby, Lootfollah Fouladi, Somayeh Najafi<sup>1</sup>, Sedigheh Asgary<sup>2</sup>, Zahra Torabi<sup>3</sup>

Islamic Azad University, Najafabad, <sup>1</sup>Acquired Immunodeficiency Research Center, <sup>2</sup>Isfahan Cardiovascular Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical, <sup>3</sup>Isfahan University of Medical Sciences, Isfahan, Iran

## Abstract

**Background:** Chromium supplementations (Cr) have been shown to exert beneficial effects in the management of type-2 diabetes. Prevalence of Cr deficiency in pre-diabetic patients is not well-understood, therefore, the aim of this study was to evaluate the extent of this prevalence.

**Materials and Methods:** In this cross-sectional descriptive study, 132 pre-diabetic patients were recruited. The participants were randomly selected from those who referred to the Shariati Hospital in Isfahan, Iran. Blood samples are collected for measurement of Cr, insulin, fasting blood sugar (FBS), and two-hour post-load plasma glucose. The body mass index (BMI) was calculated. Determination of Cr was carried out by atomic absorption spectrometry.

**Results:** Thirty-four (31.5%) patients had Cr deficiency and 74 (68.5%) patients had normal Cr. There was no significant difference between sex, age groups (<50 years and ≥50 years) and between patients with and without a family history of diabetes in both the groups. No significant differences in age, BMI, FBS or insulin were observed between two groups. In the group with a normal level of Cr, there was a significant reversed correlation between the Cr level and age, but no significant correlation existed between the Cr level and other factors in both groups.

**Conclusion:** The levels of Cr deficiency are relatively common in patients with pre-diabetes, and it is necessary to screen patients with diabetes and pre-diabetes according to the American Diabetes Association guidelines, with regard to the Cr level and action should be taken to eliminate the Cr deficiency in these patients.

**Key Words:** Chromium, insulin, pre-diabetic

### Address for correspondence:

Dr. Zahra Torabi, Isfahan University of Medical Sciences, Isfahan, Iran. E-mail: [zahratorabi194@yahoo.com](mailto:zahratorabi194@yahoo.com)

Received: 11.09.2013, Accepted: 03.02.2014

## INTRODUCTION

Diabetes mellitus, a well-known endocrine metabolic disorder, is a disease characterized by high levels of blood glucose and multiple tissue complications, resulting in nephropathy, neuropathy, and retinopathy.<sup>[1]</sup> Impaired glucose tolerance (IGT), impaired fasting glucose (IFG), and metabolic syndrome are considered precursors to type 2 diabetes mellitus.<sup>[2]</sup> The presence of IGT and IFG

Access this article online	
Quick Response Code:	Website: <a href="http://www.advbiores.net">www.advbiores.net</a>
	DOI: 10.4103/2277-9175.145737

Copyright: © 2014 Rafiei. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**How to cite this article:** Rafiei R, Habyby Z, Fouladi L, Najafi S, Asgary S, Torabi Z. Chromium level in prediction of diabetes in pre-diabetic patients. *Adv Biomed Res* 2014;3:235.

is called pre-diabetes.<sup>[3]</sup> A significant increase in the costs of long-term treatment of diabetes and its complications is felt.<sup>[4]</sup> Although pharmacotherapy can reduce the risk of type 2 diabetes,<sup>[5]</sup> their cost and potential adverse effects can be objectionable to patients who do not yet have the actual disease.<sup>[6]</sup> Intensive diet and lifestyle changes can have an important role in diabetes prevention.<sup>[7]</sup> Chromium (Cr) may help people with type 2 diabetes by improving insulin resistance and increasing the body's sensitivity to insulin.<sup>[3]</sup> Cr is considered an essential element for humans.<sup>[8]</sup> A diet lacking in this element may result in the development of diabetes mellitus.<sup>[8]</sup>

Cr is required for normal carbohydrate, protein, and lipid metabolism and its deficiency has been implicated as one of the causes of diabetes mellitus.<sup>[9,10]</sup> Most of the current knowledge on the effects of Cr on glucose homeostasis comes from clinical studies examining the effect of Cr supplementation on glucose intolerance and insulin resistance. Results from these studies have been inconclusive.<sup>[11,12]</sup>

In the present research, with regard to diabetes prevalence and a need to give special attention to its prevention and the effect of Cr on diabetes, we investigate the effect of the Cr level in the prediction of diabetes in pre-diabetic patients.

## MATERIALS AND METHODS

This cross-sectional descriptive study was designed to evaluate the effect of the Cr level on the prediction of diabetes in pre-diabetic patients. The study was conducted in 2012, and 132 pre-diabetic patients, with the mean age of  $52.8 \pm 13.3$  years, were recruited.

The participants were randomly selected from those who referred to the Shariati Hospital (Isfahan, Iran). The allocation was conducted using random numbers generated by the computer, using the participants' record numbers in the clinic. The blood tests used to measure plasma glucose and diagnose pre-diabetes included a fasting plasma glucose (FBS) test and a glucose tolerance test. Individuals with a fasting plasma glucose level between 100 and 125 mg/dl were referred to as having impaired fasting glucose. Individuals with plasma glucose levels of 140-199 mg/dl after a two-hour oral glucose tolerance test were referred to as having impaired glucose tolerance.<sup>[13]</sup> Patients were excluded if they were taking antidiabetic drugs or Cr supplementation. The demographic details were taken from the participants and their parents. This

study protocol was approved by the Medical Ethics Committee of the Islamic Azad University.

All measurements were made by a trained team of general physicians and nurses under the supervision of the same pediatrician. Anthropometric and biochemical measurements were conducted using calibrated instruments and standard protocols after participant selection. Fasting venous blood samples were collected from participants to measure the biochemical factors. Blood glucose was determined using the glucose oxidase assay and insulin levels were determined using the enzyme-linked immunosorbent assay (ELISA). A two-hour post-load plasma glucose test (2 hpp) was performed twice by venous blood sampling (in a fasting state and two hours after drinking a glucose solution) in all participants. Determination of Cr was carried out by atomic absorption spectrometry (Agilent, USA).<sup>[14]</sup> Calculation of body mass index (BMI) was done by dividing weight (in kilograms) by height (in meters squared).


Statistical analyses were performed using SPSS program (Version 15, Chicago, IL, USA). The data were presented as means with standard deviation (SD). The chi-square and Fisher tests were used for all comparisons between the qualitative data and the T test was applied to compare the quantitative data. The Pearson's correlation was used for investigation of the studied factors and Cr levels. A *P*-value of  $<0.05$  was considered to be statistically significant.

## RESULTS


In this study, 132 patients were investigated but 24 patients didn't complete their laboratory studies and were not included in the final statistical analyses. The mean and standard deviation of serum Cr was  $1.08 \pm 1.06$   $\mu\text{g/L}$ . Thirty-four (31.5%) patients had Cr deficiency and 74 (68.5%) patients had normal Cr. Table 1 illustrates that there is no significant difference between men and women, between the investigated age groups ( $<50$  years and  $\geq 50$  years), and between patients with and without a family history of diabetes, in both the groups.

According to the findings, as shown in Table 2 and Figures 1-3, no significant difference was observed in age and BMI, as well as FBS and insulin, between the two groups. Cr deficiency occurred at higher levels of FBS and BMI, but at older ages they were not statistically significant.


Findings on the correlation between Cr and the biochemical factors of blood in the normal Cr and the



**Figure 1:** The Cr levels (µg/L) based on sex in two normal Cr and Cr-deficiency groups



**Figure 2:** The Cr levels (µg/L) based on age in two normal Cr and Cr-deficiency groups



**Figure 3:** The Cr levels (µg/L) based on family history of diabetes in two normal Cr and Cr-deficiency groups

Cr-deficiency groups are summarized in Table 3 and Figures 4-7. In the group with a normal level of Cr, there was a significant reversed correlation between the Cr level and age, but no significant correlation existed between the Cr level and other factors in both groups.

## DISCUSSION

Chromium is believed to be an insulin-sensitizing agent and may facilitate insulin attachment to the insulin receptor.<sup>[15]</sup> Cr may improve insulin sensitivity by activating the insulin receptor tyrosine kinase, an effect that has been shown in rats,<sup>[16]</sup> and also by inhibiting phosphotyrosine phosphatase, which is a rat homolog of human tyrosine phosphatase, which inactivates the activated (phosphorylated) insulin

**Table 1:** The Cr levels (µg/L) based on sex, age, and family history of diabetes in two normal Cr and Cr-deficiency groups

Group	Anthropometric variable	Mean ± SD	P-value
Normal Cr	Female	1.56±1.01	0.509
	Male	1.39±1.1	
Cr deficiency	Female	0.15±0.11	0.349
	Male	0.19±0.12	
Normal Cr	<50 years	1.79±1.32	0.083
	≥50 years	1.31±0.75	
Cr deficiency	<50 years	0.15±0.10	0.952
	≥50 years	0.17±0.12	
Normal Cr	Without a family history of diabetes	1.47±0.91	0.733
	With a family history of diabetes	1.55±1.19	
Cr deficiency	Without a family history of diabetes	0.17±0.12	0.789
	With a family history of diabetes	0.16±0.11	

The results are expressed as mean values ± standard deviation (SD), P-values are significant P < 0.05

**Table 2:** Biochemical and anthropometric variables in two normal Cr and Cr-deficiency groups

Variable	Group	mean ± SD	P-value
Age	Normal Cr	51.54±13.92	0.153
	Cr deficiency	55.47±11.37	
FBS	Normal Cr	107.57±8.51	0.626
	Cr deficiency	108.47±9.76	
BMI	Normal Cr	29.62±4.62	0.189
	Cr deficiency	28.18±6.14	
Insulin	Normal Cr	15.01±24.55	0.981
	Cr deficiency	14.86±21.76	

The results are expressed as mean values ± standard deviation (SD), P-values are significant P < 0.05

**Table 3:** Correlation between Cr level and biochemical and anthropometric variables in two normal Cr and CrMean of age in two-deficiency groups

Variable	Group	Pearson's correlation	P-value
Age	Normal Cr	-0.276	0.017
	Cr deficiency	0.102	
Two-hour post load plasma glucose	Normal Cr	0.038	0.774
	Cr deficiency	-0.218	
BMI	Normal Cr	0.224	0.055
	Cr deficiency	-0.071	
FBS	Normal Cr	-0.069	0.558
	Cr deficiency	0.276	
Insulin	Normal Cr	-0.106	0.509
	Cr deficiency	-0.268	

P-values are significant P < 0.05

receptor. Thus, Cr may promote phosphorylation, and hence, the activation of the insulin receptor, leading to an improvement in insulin sensitivity.<sup>[16]</sup>

One of the reviews has identified 16 studies on Cr supplementation in patients with type 2 diabetes.




Figure 4: Mean of age in two normal Cr and Cr-deficiency groups




Figure 6: Mean of BMI in two normal Cr and Cr-deficiency groups




Figure 5: Mean of FBS in two normal Cr and Cr-deficiency groups




Figure 7: Mean of insulin in two normal Cr and Cr-deficiency groups

Of these studies, 13 show a significant improvement in glucose, insulin, and lipid concentrations. These studies include a total of 502 patients. There are three studies, with a total of 55 subjects, in which no effect has been found.<sup>[17]</sup> Cr has a positive effect on glucose tolerance and diabetes, but all the studies do not involve these effects for a number of reasons. First of all, human studies include subjects of diverse genetic and nutritional backgrounds, living in environments of varying degrees of stress, all of which may affect Cr metabolism.<sup>[18]</sup> Varying results of supplemental Cr may also be due to the diet, selection of subjects, the duration of the study, as also the amount and type of supplemental Cr. In addition, response to Cr is related to the degree of glucose intolerance. Subjects with better glucose tolerance, who do not need additional Cr, do not respond to supplemental Cr.<sup>[19]</sup>

The study by Tripathy *et al.*<sup>[20]</sup> was conducted on 30 type-2 diabetic subjects and 30 age-matched control subjects within the age range of 40-55 years. Glucose and Cr were analyzed in the fasting serum of all the subjects. Diabetic subjects had significantly lower serum concentrations of Cr as compared to the controls and exhibited a significant increase in serum glucose levels. The serum levels of glucose were found to be negatively correlated to the serum levels of Cr in diabetic patients. The lower serum levels of Cr in diabetics compared to control patients could be due to the poor glycemic control.<sup>[20]</sup> Cr supplementation had variable effects on the body weight of patients with diabetes.<sup>[21,22]</sup> One study of patients with diabetes using Cr indicated no significant effects on either body weight or BMI,<sup>[21]</sup> while of the eight double-blind, placebo-

controlled trials in individuals without diabetes, Cr supplementation showed a decrease in weight and fat in three larger studies.<sup>[22]</sup> Ding *et al.*<sup>[23]</sup> revealed that serum Cr was similar in women and men, while in the results of the Ravina *et al.*<sup>[24]</sup> study, the blood Cr level was lower in women than in men. In the present study, the number of women with Cr deficiency was more than men, which was consistent with the Ding *et al.*<sup>[23]</sup> study. The difference in the results could be due to differences in nutrition and geographic area. Duncan *et al.*<sup>[25]</sup> concluded that increasing age and family history of diabetes led to insulin resistance in diabetic patients, due to reduction in blood levels of Cr. In our study, in patients with a deficiency of Cr, the number of patients  $\geq 50$  years old and with a family history of diabetes was more than the others, however, it was not significant. Based on the results achieved from this study, the number of pre-diabetic patients with deficiency of Cr than patients with normal level of Cr was not negligible, and it could be seen as a warning for the onset of diabetes.

According to the effect of varying factors, including nutrition, race and environment played a role in diabetes. This was suggested, and further studies should be conducted, with larger sample sizes. Risk factors for diabetes investigate in the future also.

## CONCLUSION

With regard to the Cr deficiency in pre-diabetes patients and the effect of this element on blood glucose and insulin sensitivity, screening for pre-diabetic should be done (according to the American Diabetes

Association) and the levels of Cr be investigated in these patients in order to perform the appropriate action for the removal of Cr deficiency and prediction of diabetes.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge the authorities at the Islamic Azad University, Najafabad branch, for their financial support for this study.

## REFERENCES

1. Sicree R, Shaw JE, Zimmet PZ. The global burden of diabetes. In: Gan D, editor. *Diabetes Atlas*. 2<sup>nd</sup> ed. Brussels, Belgium: International Diabetes Federation;2003. p. 15-71.
2. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2009;32(Suppl 1):S62-7.
3. Ali A, Ma Y, Reynolds J, Wise JP Sr, Inzucchi SE, Katz DL. Chromium effects on glucose tolerance and insulin sensitivity in persons at risk for diabetes mellitus. *Endocr Pract* 2011;17:16-25.
4. Parkin CG, Davidson JA. Value of self-monitoring blood glucose pattern analysis in improving diabetes outcomes. *J Diabetes Sci Technol* 2009;3:500-8.
5. Padwal R, Majumdar SR, Johnson JA, Varney J, McAlister FA. A systematic review of drug therapy to delay or prevent type 2 diabetes. *Diabetes Care* 2005;28:736-44.
6. Walker EA, Molitch M, Kramer MK, Kahn S, Ma Y, Edelstein S, *et al.* Adherence to preventive medications: Predictors and outcomes in the Diabetes Prevention Program. *Diabetes Care* 2006;29:1997-2002.
7. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, *et al.*; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393-403.
8. Afridi HI, Kazi TG, Brabazon D, Naher S, Talpur FN. Comparative metal distribution in scalp hair of Pakistani and Irish referents and diabetes mellitus patients. *Clin Chim Acta* 2013;415:207-14.
9. Wang YQ, Yao MH. Effects of chromium picolinate on glucose uptake in insulin-resistant 3T3-L1 adipocytes involve activation of p38 MAPK. *J Nutr Biochem* 2009;20:982-91.
10. Kwon MJ, Chung HS, Yoon CS, Ko JH, Jun HJ, Kim TK, *et al.* The effect of chromium on rat insulinoma cells in high glucose conditions. *Life Sci* 2010;87:401-4.
11. Balk EM, Tatsioni A, Lichtenstein AH, Lau J, Pittas AG. Effect of chromium supplementation on glucose metabolism and lipids: A systematic review of randomized controlled trials. *Diabetes Care* 2007;30:2154-63.
12. Vincent JB. *The Nutritional Biochemistry of Chromium (III)*. Amsterdam, Boston: Elsevier; 2007.
13. Phillips LS, Weintraub WS, Ziemer DC, Kolm P, Foster JK, Vaccarino V, *et al.* All pre-diabetes is not the same: Metabolic and vascular risks of impaired fasting glucose at 100 versus 110 mg/dl: The screening for impaired glucose tolerance study 1 (SIGT 1). *Diabetes Care* 2006;29:1405-7.
14. Nawrocka A, Szkoda J. Determination of chromium in biological material by electrothermal atomic absorption spectrometry method. *Bull Vet Inst Pulawy* 2012;56:585-9.
15. Mertz W, Toepfer EW, Roginski EE, Polansky MM. Present knowledge of the role of chromium. *Fed Proc* 1974;33:2275-80.
16. Davis CM, Vincent JB. Chromium oligopeptide activates insulin receptor tyrosine kinase activity. *Biochemistry* 1997;36:4382-5.
17. Gunton JE, Hams G, Hitchman R, McElduff A. Serum chromium does not predict glucose tolerance in late pregnancy. *Am J Clin Nutr* 2001;73:99-104.
18. Anderson RA. Stress effects on chromium nutrition of humans and farm animals. In: Anonymous, editor. *Proceedings of Alltech's Tenth Symposium on Biotechnology in the Feed Industry*. Nottingham, England: University Press;1994. p. 267-74.
19. Anderson RA. Recent advances in the clinical and biochemical effects of chromium deficiency. In: Prasad AS, editor. *Essential and Toxic Trace Elements in Human Health and Disease*. New York: Wiley Liss;1993. p. 221-34.
20. Tripathy S, Sumathi S, Raj GB. Minerals nutritional status of type 2 diabetic subjects. *Int J Diab Dev Countries* 2004;24:27-8.
21. Cefalu WT, Rood J, Pinsonat P, Qin J, Sereda O, Levitan L, *et al.* Characterization of the metabolic and physiologic response to chromium supplementation in subjects with type 2 diabetes mellitus. *Metabolism* 2010;59:755-62.
22. Campbell WW, Joseph LJ, Anderson RA, Davey SL, Hinton J, Evans WJ. Effects of resistive training and chromium picolinate on body composition and skeletal muscle size in older women. *Int J Sport Nutr Exerc Metab* 2002;12:125-35.
23. Ding W, Chai Z, Duan P, Feng W, Qian Q. Serum and urine chromium concentrations in elderly diabetics. *Biol Trace Elem Res* 1998;63:231-7.
24. Ravina A, Slezack L. Chromium in the treatment of clinical diabetes mellitus. *IMAJ* 1993;125:142-5, 191.
25. Duncan MG. The effects of nutritional supplements on the treatment of depression, diabetes, and hypercholesterolemia in the renal patient. *J Ren Nutr* 1999;9:58-62.

Source of Support: Nil, Conflict of Interest: None declared.