

Evaluation of the effect of *Cyperus rotundus* L. in scopolamine-induced learning deficit in mice

Mohammed Rabbani, Alireza Ghannadi¹, Nahid Malekian

Department of Pharmacology, School of Pharmacy and Pharmaceutical Sciences and Isfahan Pharmaceutical Sciences Research Center,

¹Department of Pharmacognosy, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract **Background:** *Cyperus rotundus* L. was used in traditional Iranian medicine to treat memory and cognition disorders. The present study was aimed at investigating the effect of the extract and essential oil of *C. rotundus* on memory dysfunction.

Materials and Methods: Cognition was evaluated using the object recognition task that was composed of a square wooden open field box with different shape objects. The test was consisted of three sections: 15 min exploration, first trial for 12 min and second one for 5 min. In the second trial the difference in exploration between a previously seen object and novel one, was considered as an index of memory performance (recognition index). Memory deficit was induced by scopolamine (0.5 mg/kg) before injection of plant extracts and essential oil.

Results: Rivastigmine at 0.6 mg/kg reversed the scopolamine induced memory dysfunction in mice ($P < 0.05$). On the contrary, neither the hydroalcoholic extracts (100, 200, 400 mg/kg) nor the polyphenolic extract (50, 100, 200 mg/kg) and essential oil (10, 20, 40 mg/kg) of *C. rotundus* produced significant improvement of memory dysfunction. The fact that rivastigmine reversed the scopolamine-induced memory dysfunction confirms the validity of this memory paradigm.

Conclusion: Using the current method of the memory evaluation, none of the tested doses of the plant extract or essential oil changed the memory status of the animals, indicating either a lack of effective ingredient or unsuitable method for evaluation.

Key Words: *Cyperus rotundus*, memory dysfunction, object recognition, scopolamine

Address for correspondence:

Prof. Mohammed Rabbani, Department of Pharmacology, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran. E-mail: rabanim@yahoo.com

Received: 03.10.2012, **Accepted:** 26.02.2013

INTRODUCTION

Memory is one of the most important and complicated behavioral process that study of its physiologic and

anatomic principles can help prevents memory loss in different conditions. The cause for most memory deficient cases (such as Alzheimer, Parkinson, amnesia, and dementia) is still unknown. Accumulation of amyloid proteins that causes the degeneration of cholinergic neurons, which in turn results in cholinergic deficiency and synapse loss is the best know hypothesis that has been put forward.^[1] Based on these studies, acetylcholinesterase (AChE) inhibitors have been used to treat patients with cognitive impairment.^[2] Classical AChE inhibitor drugs however, provide short-term alleviation of symptoms without decelerating the progression of

Access this article online	
Quick Response Code:	Website: www.advbiores.net
	DOI: 10.4103/2277-9175.143293

Copyright: © 2014 Rabbani. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article: Rabbani M, Ghannadi A, Malekian N. Evaluation of the effect of *Cyperus rotundus* L. in scopolamine-induced learning deficit in mice. Adv Biomed Res 2014;3:217.

the disease.^[3] Therefore, the development of new drugs that somehow counteract the progression of the disease would be highly desirable for treating such patients. Mental stimulation, exercise and balanced diet have been suggested as other useful ways to delay symptoms.^[4,5]

Cyperus rotundus L. (family *Cyperaceae*), also known as purple nutsedge or nutgrass, is a common perennial weed and scaly creeping rhizome. The tubers are externally brown in color and reddish white inside with a characteristic odor.^[6] The rhizome of *C. rotundus* is a well-known functional food and traditional herbal medicine in Asia.^[7] In traditional herbal medicine in Iran, the rhizome of *C. rotundus* is used for memory, dermatological and psychological disorders.^[8,9] The rhizome of this plant has been reported to show antioxidant and free radical scavenging activities, these effects play a major role in protection of neurodegenerative disorders.^[10] In addition to these, the plant has also antifungal, anti-inflammatory, anti-diabetic, cyto protective, anti-microbial, antioxidants, and cytotoxic properties.^[6,11,12] Considering the wide-spread central action of the plant and in order to verify its traditional use as a memory enhancer, the present study was undertaken to determine the effect of *C. rotundus* on memory performance after scopolamine-induced neuronal damage.^[4] Memory is assessed by the novel object recognition task that originally developed by Ennaceur and Delacour in 1988 and it is based on natural tendency of rodent to explore a novel object more than familiar one.^[13]

MATERIALS AND METHODS

Animals

Male mice (8-12 weeks, Pasteur Institute, Tehran, Iran) with average weight of 18-35 g were housed in cage of six in the experiment place a day before study. Tap water and standard food pellets were available *ad libitum*. All the studies were performed between 8 AM and 12 AM in a special noise-free room with controlled illumination. There were at least six mice in each group.

Drug treatments

As a control group mice was treated with saline, and for inducing memory dysfunction scopolamine was injected at doses 0.5 mg/kg and 1 mg/kg (the dose 1 mg/kg was administered only for negative control group). For positive control group rivastigmine was administered 0.6 mg/kg and the test groups were injected with hydroalcoholic extracts (100, 200, 400 mg/kg), polyphenolic extracts (50, 100, 200 mg/kg) and essential oil (10, 20, 40 mg/kg) 5 min

after scopolamine injection. Then the animals were kept in their home cage for 25 min and finally the mice were placed in the test box for starting T1 [Figure 1].

Sample preparation

Scopolamine sulphate (Daru pakhsh, Iran) was diluted in normal saline (0.9%). Rivastigmine (Modava, Tehran, Iran) and was prepared in normal saline 0.9% and tween 80 (1% v/v) containing 1% (w/v) methylcellulose.^[14] All polyphenolic extracts were diluted with water, hydroalcoholic extracts and essential oil were diluted with water and tween 80 (1% v/v). Control animals received either normal saline or vehicle. All of the drugs were injected intraperitoneally (i.p.).

Tubers of *C. rotundus* were collected on October 2010 from plants growing wild in Baharan homestead in Kelishad area in Isfahan province in Iran at altitude of ca 1650 m. The plant was authenticated by Dr. Lili Ghaem maghami (Biology Department, Isfahan University, Iran), as *C. rotundus* L. The plant herbarium was deposited in the School of Pharmacy, at Isfahan University of Medical Sciences with code number of 2262.

For preparation of hydroalcoholic extract, air-dried and powdered tubers of plant (300 g) were macerated with Ethanol: H₂O (7:3) for 48 h. The extract was then shaken, filtered and evaporated in a rotating evaporator.^[15] For preparation of the polyphenolic extract, air-dried and powdered tubers of the plant (300 g) were extracted in two steps: firstly with Ethanol: H₂O (9:1) and secondary with Ethanol: H₂O (1:1) For 48 h. The two extracts were combined and evaporated to about 1/3 of the original volume. The resultant aqueous extract was cleared by extraction in a separating funnel with chloroform and then evaporated to dryness under reduced pressure in a rotating evaporator.^[16] The essential oil was isolated by hydrodistillation of the air-dried powdered tubers of the plant for 3-4 h based on our previous studies. Essential oil sample was homogenized and dried over anhydrous sodium sulfate and stored in fridge.^[15]




Figure 1: Effect of rivastigmine on scopolamine induced memory deficit. The first injection was made 30 min prior to test and consisted of saline for control groups and scopolamine for others. This was followed 5 min later by a second injection (rivastigmine, extracts, and essential oil). The animals were kept in the home cage for 30 min before starting the test

Object recognition test

Novel object recognition task is a relatively sensitive procedure for evaluating compounds effects on cognition activity. The apparatus was consisting of square wooden cage (35 × 35 × 40 cm) with the black walls, which located in noise free room. The day before the test, each animal was submitted to a habituation session in the open field and allowed to freely explore the arena in the presence of two objects for at least 15 min on experimental day, animals were submitted to two trials spaced by an intertribal interval (15 min). During the first trial (acquisition trial, T1), animals were placed in the arena containing two identical objects for an amount of time necessary to explore the objects for 20 s. Mice that did not explore the objects for 20 s within the 12-min period was exclude from the experiments. Exploration is defined as the animal directing the nose within 2 cm of the object while looking at, sniffing, or touching it. For the second trial (test trial, T2), one of the objects presented in the first trial was replaced by a new object, animals were placed back in the arena for 5 min and total time spent in exploration of each object was determined. Animals behavior were recorded by using an ordinary web camera mounted above the experimental apparatus, records were analyzed later.^[17]

Data processing and statistical analysis

The parameters were measured, time required to achieve 20 s of object exploration on T1, time spent in active exploration of the familiar (F) or novel (N) object on T2. Recognition index (RI) was defined using the following formula: $(N - F)/(N + F) \times 100$. All the procedures were recorded using a camera, which was placed above the experimental apparatus.^[17] T1 was recorded for each mouse, RI was calculated, and the data were analyzed using two-way ANOVA and student *t*-test, $P < 0.05$ was considered as significant result. Results were expressed as mean + SEM. Sigma stat software was used for statistical analyses.

RESULTS

The essential oil of *C. rotundus* was a yellow liquid bearing the characteristic pungent and aromatic odor of *Cyperaceae* plants. The essential oil content of the plant was 0.2% (w/w) yield. Evaporation and solvent removal of hydroalcoholic extract gave a semisolid mass with a yield of 19.0% (w/w) and yield of polyphenolic was 8.3% (w/w).

Effect of scopolamine on RI

In this study, mice received 0.5 mg/kg and 1 mg/kg of scopolamine were significantly different with saline group [Figure 2]. Figure 2 shows the RI in the second trial of object recognition task after scopolamine.




Figure 2: Effect of different doses of scopolamine on memory performance (expressed as recognition index) and effect of rivastigmine (0.6 mg/kg) on reversal of memory deficit induced by scopolamine. Results are expressed as mean + SEM ($n = 6$). * $P < 0.05$ compared with saline. $^{\$}P < 0.05$ compared with scopolamine

The ANOVA revealed a significant treatment effect ($P < 0.05$). The RI score was decreased in the scopolamine condition compared to the saline condition, as shown by the *post-hoc* analysis.

Effect of rivastigmine on RI

Based on previous studies 0.6 mg/kg of rivastigmine was used to test reversal of memory impairment by scopolamine. At this dose the memory was reversed to control level. There was no significant difference between rivastigmine and saline.

Figure 2 shows the RI in the second trial of object recognition task after scopolamine and rivastigmine administration. The ANOVA revealed a significant treatment effect ($P < 0.05$). The RI score was increased in the scopolamine induced memory deficit compared to the saline condition.

Effect of extracts and essential oil in scopolamine-induced memory deficit

The RI of mice received 50, 100, and 200 mg/kg of polyphenolic extract, 100, 200, and 400 mg/kg hydroalcoholic extracts and 10, 20, and 40 mg/kg essential oil of *C. rotundus* were not significantly different with saline [Figures 3-5].

DISCUSSION

Herbal drugs are proved to be of great commercial and pharmacological importance. Traditional herbal medicine forms an important part the health-care system. Plants have been used for centuries as remedies for human diseases.^[18] *C. rotundus*, a member of *Cyperaceae* family is grown throughout the world from tropics to temperate regions. In Iranian traditional medicine *C. rotundus* has long been used




Figure 3: Effect of scopolamine on memory performance (expressed as recognition index) and effect of different doses of *Cyperus rotundus* hydroalcoholic extract on reverse of memory deficit induced by scopolamine. Results are expressed as mean + SEM ($n = 6$). * $P < 0.05$ compared with saline




Figure 4: Effect of scopolamine on memory performance (expressed as recognition index) and effect of different doses of *Cyperus rotundus* polyphenolic extract on reverse of memory deficit induced by scopolamine. Results are expressed as mean + SEM ($n = 6$). * $P < 0.05$ compared with saline




Figure 5: Effect of scopolamine on memory performance (expressed as recognition index) and effect of different doses of *Cyperus rotundus* essential oil on reverse of memory deficit induced by scopolamine. Results are expressed as mean + SEM ($n = 6$). * $P < 0.05$ compared with saline

to improve memory.^[8] Despite its wide-spread use as a memory enhancer, no pharmacological data is available to support such claim. Keeping in view its suitability as food ingredient, the current investigation was designed to investigate the effect of *C. rotundus* on scopolamine-induced memory dysfunction in mice using object recognition task.

In the current work, scopolamine, a competitive muscarinic antagonist reduced the RI by 66%. Amnesic properties of scopolamine, have long been known in humans and animals, and in fact, scopolamine amnesia has been proposed as a pharmacological model for human dementia and for Alzheimer's disease.^[19] The range of memory dysfunction in the similar studies that have used scopolamine was 70%.^[17]

Reversal of scopolamine induced memory deficit has been used as an initial screening method for identification of therapeutic candidate for cognitive disorders.^[19] Rivastigmine, an acetylcholine esterase inhibitor is routinely used as a control drug for testing the validity of the method.^[20,21] In the present work, the scopolamine-induced memory deficit was reversed to control value by 0.6 mg/kg of rivastigmine. Although, no other study has used rivastigmine as a control in object recognition test, in other memory paradigm model, the result of rivastigmine is very similar to that of ours reverse scopolamine induced in some dose.^[14]

The plant extract and essence used in varying doses in the present study failed to improve the impaired memory by scopolamine. This could be due to several factors, such as the animal model or the underlying mechanism that was studied here. Several animal models are available for studying such function that include, radial maze, water maze, and object recognition paradigm.^[22] It is well-established that in object recognition model of memory both the short- and long-term memory is studied.^[23] Since in the present experimental setting the whole procedure takes 24 h to complete, we are only looking at the short term memory.^[23] It is possible that for the effective action, the plant extract requires to be present over a longer period of time. As far as the memory performance model is concerned, this model is considered as a simple and reliable method that is fast and reproducible.^[17] However, there are some studies that shows some drugs are effective in one model without a significant effects on other model.^[24]

Since in several Iranian traditional medicine resources and literatures the *C. rotundus* is recommended to be used in combination with other plants such as *Crocus sativus*, *Zingiber officianale*, and *Boswellia papyrifera*^[25] it is possible that the effect that

is produced to be due to the presence of other plant ingredients like *B. papyrifera*. The memory improvement property of *B. papyrifera* has already been shown by number of studies.^[26]

As mentioned above, in the object recognition model we used scopolamine to impair the animal's memory. Therefore, any drugs that could reverse the action of scopolamine such as rivastigmine could produce effective action and drugs that are not acting through this remain ineffective. One other possibility for not seeing significant memory improvement effect by the plant extract in this study could be due to lack of cholinergic action. It is well-known that *C. rotundus* has antioxidant properties.^[10,27] It is therefore, possible that the memory enhancement by *C. rotundus* to be due to the presence of such components in the extract of the plants. Different experimental memory set up is required to evaluate the antioxidant properties of the drugs.^[28]

CONCLUSION

In summary, the present study shows that the *C. rotundus* extracts or essence do not provide any memory impairment in mice. Lack of active ingredient in the plant seems to be the major reason for not observing any effects as this preclinical model. Further studies are underway to test the combination of this plant and other known so-called memory enhancer plants.

REFERENCES

1. Terry RD, Masliah E, Salmon DP, Butters N, DeTeresa R, Hill R, et al. Physical basis of cognitive alterations in Alzheimer's disease: Synapse loss is the major correlate of cognitive impairment. *Ann Neurol* 1991;30:572-80.
2. Cummings JL, Mackell J, Kaufer D. Behavioral effects of current Alzheimer's disease treatments: A descriptive review. *Alzheimers Dement* 2008;4:49-60.
3. Karran E, Mercken M, De Strooper B. The amyloid cascade hypothesis for Alzheimer's disease: An appraisal for the development of therapeutics. *Nat Rev Drug Discov* 2011;10:698-712.
4. Babic T. The cholinergic hypothesis of Alzheimer's disease: A review of progress. *J Neurol Neurosurg Psychiatry* 1999;67:558.
5. Aarsland D, Cummings JL, Larsen JP. Neuropsychiatric differences between Parkinson's disease with dementia and Alzheimer's disease. *Int J Geriatr Psychiatry* 2001;16:184-91.
6. Lawal OA, Oyediji AO. Chemical composition of the essential oils of *Cyperus rotundus* L. from South Africa. *Molecules* 2009;14:2909-17.
7. Lee CH, Hwang DS, Kim HG, Oh H, Park H, Cho JH, et al. Protective effect of *Cyperus rotundus* rhizoma against 6-hydroxydopamine-induced neuronal damage. *J Med Food* 2010;13:564-71.
8. Rhazes MZ, Al-Hawi. Vol. 21. Tehran: Publications of Iranian Academy of Medical Sciences; 2005. p. 16.
9. Zargari A. Medicinal plants. Vol. 4. Tehran: Tehran University Press; 1990. p. 681-2.
10. Yazdanparast R, Ardestani A. *In vitro* antioxidant and free radical scavenging activity of *Cyperus rotundus*. *J Med Food* 2007;10:667-74.
11. El-Gohary HM. 14-Study of essential oils of the tubers of *Cyperus rotundus* L. and *Cyperus alopecuroides* rottb. *Bull Fac Pharm Cairo Univ* 2004;42:157.
12. Nassar MI, Abdel-Razik AF, El-Khrisy Eel-D, Dawidar AA, Bystrom A, Mabry TJ. A benzoquinone and flavonoids from *Cyperus alopecuroides*. *Phytochemistry* 2002;60:385-7.
13. Ennaceur A, Delacour J. A new one-trial test for neurobiological studies of memory in rats. 1: Behavioral data. *Behav Brain Res* 1988;31:47-59.
14. Deiana S, Harrington CR, Wischik CM, Riedel G. Methylthionium chloride reverses cognitive deficits induced by scopolamine: Comparison with rivastigmine. *Psychopharmacology (Berl)* 2009;202:53-65.
15. Hajhashemi V, Ghannadi A, Hajiloo M. Analgesic and anti-inflammatory effects of *Rosa damascena* hydroalcoholic extract and its essential oil in animal models. *IJPR* 2010;9:163-8.
16. Jafarian A, Ghannadi A, Elyasi A. The effects of *Allium hirtifolium* Boiss. On cell-mediated immune response in mice. *IJPR* 2010;2:51-5.
17. Bertaina-Anglade V, Enjuanes E, Morillon D, Drieu la Rochelle C. The object recognition task in rats and mice: A simple and rapid model in safety pharmacology to detect amnesic properties of a new chemical entity. *J Pharmacol Toxicol Methods* 2006;54:99-105.
18. Azaizeh H, Fulder S, Khalil K, Said O. Ethnobotanical knowledge of local Arab practitioners in the Middle Eastern region. *Fitoterapia* 2003;74:98-108.
19. Bartus RT, Dean RL, Pontecorvo MJ, Flicker C. The cholinergic hypothesis: A historical overview, current perspective, and future directions. *Ann N Y Acad Sci* 1985;444:332-58.
20. de Bruin N, Pouzet B. Beneficial effects of galantamine on performance in the object recognition task in Swiss mice: Deficits induced by scopolamine and by prolonging the retention interval. *Pharmacol Biochem Behav* 2006;85:253-60.
21. Sambeth A, Riedel WJ, Smits LT, Blokland A. Cholinergic drugs affect novel object recognition in rats: Relation with hippocampal EEG? *Eur J Pharmacol* 2007;572:151-9.
22. Dere E, Huston JP, De Souza Silva MA. The pharmacology, neuroanatomy and neurogenetics of one-trial object recognition in rodents. *Neurosci Biobehav Rev* 2007;31:673-704.
23. Abush H, Akirav I. Short- and long-term cognitive effects of chronic cannabinoids administration in late-adolescence rats. *PLoS One* 2012;7:e31731.
24. Park SJ, Jung JM, Lee HE, Lee YW, Kim DH, Kim JM, et al. The memory ameliorating effects of INM-176, an ethanolic extract of *Angelica gigas*, against scopolamine- or A β (1-42)-induced cognitive dysfunction in mice. *J Ethnopharmacol* 2012;143:611-20.
25. Vasudevan M, Parle M. Evidence of anti-demential potential of abana: an Indian ayurvedic poly-herbal formulation in rats. *Pharmacologyonline* 2008. p. 439-54.
26. Farshchi A, Ghiasi G, Farshchi S, Malek Khatabi P. Effects of *boswellia papyrifera* gum extract on learning and memory in mice and rats. *Ir J Basic Med Sci* 2010;13:9-15.
27. Khalili M, Kiasalari Z, Roghani M, Azizi Y. Anticonvulsant and antioxidant effect of hydroalcoholic extract of *Cyperus rotundus* rhizome on pentylenetetrazole-induced kindling model in male mice. *J Med Plant Res* 2011;5:868.
28. Farr SA, Poon HF, Dogrukol-Ak D, Drake J, Banks WA, Eyerman E, et al. The antioxidants alpha-lipoic acid and N-acetylcysteine reverse memory impairment and brain oxidative stress in aged SAMP8 mice. *J Neurochem* 2003;84:1173-83.

Source of Support: This work was financially supported by research council of the Isfahan University of Medical Sciences, project number 188047, Isfahan, Iran. **Conflict of Interest:** None declared.