

Case Report

Supraglottic paraganglioma originated from superior laryngeal nerve

Ali Akbar Beigi, Afrooz Eshaghian¹

Departments of Surgery and ¹Otorhinolaryngology, Head and Neck Surgery, Al-Zahra Hospital, Isfahan University of Medical Science, Isfahan, Iran

Abstract

Neurogenic tumors, especially paraganglioma of larynx, are rare. In this article, we present a 64-year-old woman who complained of intermittent dysphagia to solid foods. Further evaluation revealed a supraglottic paraganglioma and she was treated successfully by total excision of tumor.

Key Words: Carotid body tumor, laryngeal nerves, paraganglioma

Address for correspondence:

Dr. Afrooz Eshaghian, Department of Otorhinolaryngology, Sofe St, Al-Zahra Hospital, Isfahan University of Medical Science, Isfahan, Iran.

E-mail: eshaghian@edc.mui.ac.ir

Received: 03.04.2012, Accepted: 02.08.2012

INTRODUCTION

Neuroendocrine tumors of larynx are divided into epithelial (carcinomas) and neural type lesions (paragangliomas).^[1] These types can overlap clinically and histologically. So, recently, they have been categorized into five tumor types: Typical carcinoid, atypical carcinoid, small cell neuroendocrine carcinoma, large cell neuroendocrine carcinoma, and paraganglioma.^[2]

Paraganglioma is among neuroendocrine tumors and accounts for 0.012% of all tumors.^[3] This tumor specially originates from parasympathetic nervous system and is predominant in females. The most common site of laryngeal paraganglioma is supraglottis; and it

presents with dysphonia, hemoptysia, and airway compromise.^[4]

Computed tomography (CT) and/or magnetic resonance imaging (MRI) can reveal vascularity of the lesion. Preoperative angiography and possible embolization is helpful before surgical intervention. Immunohistochemical staining establishes diagnosis of paraganglioma from other neuroendocrine tumors.^[5]

In this article, we describe a very rare presentation of paraganglioma in supraglottis.

CASE REPORT

A 64-year-old woman presented to our clinic with intermittent dysphagia to solid foods and dyspnea on effort from 2 months. She had history of goiter and gastroesophageal reflux; and was treated with Levothyroxin and Omeprazole.

Indirect laryngoscopy revealed a left mass lesion at the level of supraglottis. Direct laryngoscopy and biopsy of mass was done. Biopsy revealed polypoid tissue lined by stratified squamous epithelium; acanthosis

Access this article online	
Quick Response Code:	Website: www.advbiores.net
	DOI: 10.4103/2277-9175.125754

Copyright: © 2014 Beigi. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article: Beigi AA, Eshaghian A. Supraglottic paraganglioma originated from superior laryngeal nerve. *Adv Biomed Res* 2014;3:46.

was seen; Exocytosis of lymphocytes and neutrophils in surface epithelium was present. The underlying stroma was edematous and had congested vessels. No evidence of malignancy was seen.

In neck sonography, a solid hypoechoic hypervascular mass (34 × 27 mm) was seen in the left paramedian, between left common carotid artery and larynx. There were two hypoechoic thyroid nodules (7 mm diameter in inferior pole and 4 mm in superior pole).

Spiral neck CT scan with contrast revealed a large hypervascular mass which extended from epiglottis to the left vocal cord. It filled the left pyriform sinus, and narrowed airway tract with erosion to superior part of thyroid cartilage [Figure 1].

Biopsy from intraluminal space of hypopharynx was not diagnostic, so excision biopsy of mass was done.

With classic neck incision and exploration of carotid artery and vagus nerve, we found the tumor originated from superior laryngeal nerve, extending to supraglottic area. With preservation of superior laryngeal nerve, the tumor was dissected from supraglottic mucosa and excised completely. Because of edema of airway, prophylactic tracheostomy was done. Three days after the surgery, tracheostomy removed. Microscopic examination revealed the tumor as carotid body tumor (paraganglioma).

After surgery, the patient suffered from dysphonia as a result of superior laryngeal nerve paresis that improved 3 months later.

DISCUSSION




Figure 1: Axial CT scan of neck revealed hyperdense left pyriform sinus mass and narrowed airway tract, with erosion to superior part of thyroid cartilage

Only 10% of paragangliomas have extra-adrenal localization and only about 0.33% of them localize in head and neck. Typically they are benign tumors, and only 19% of cases may be potentially malignant. Neck paragangliomas are located close to carotid artery bifurcation, jugular bulb, and along the course of vagus nerve. They are very rare in larynx.^[3]

This tumor presents by different manifestations; mainly there is a slow-growing, painless, asymptomatic mass with compressive effect on the surrounding structures. Auscultation may reveal a carotid pulse, bruit, or thrill.^[6,7] It is more likely in females of age between 50 and 70 years.^[8]

We present a supraglottic involvement of carotid body tumor, which is a very rare site for this tumor. Over 83 patients have been described in the literature.^[9]

Diagnosis of this tumor is made by digital subtraction angiography (DSA) or Color Doppler sonography very well.^[6]

This benign tumor is treated by complete surgical excision. As the risk of bleeding is high in this tumor, some authors suggest the external approach for tumor removal;^[10] for example, lateral thyrotomy, lateral pharyngotomy, or laryngofissure technique.^[11] If the tumor size is small, it can be excised with direct laryngoscopy and endoscopically.^[12] Laser-assisted microlaryngoscopy is described for excision of large supraglottic paraganglioma as well.^[13]

In conclusion, in patient with hypervascular neck mass, carotid body tumor should be considered, although supraglottis is its unusual location, and fine-needle aspiration should be done very cautiously to prevent massive hemorrhage. High clinical suspicion is mandatory to diagnose the tumor, even if nondiagnostic biopsy has been done.

REFERENCES

1. Ferlito A, Rosai J. Terminology and classification of neuroendocrine neoplasms of the larynx. *ORL J Otorhinolaryngol Relat Spec* 1991;53:185-7.
2. Lewis JS Jr, Ferlito A, Gnepp DR, Rinaldo A, Devaney KO, Silver CE, et al. International Head and Neck Scientific Group. Terminology and classification of neuroendocrine neoplasms of the larynx. *Laryngoscope* 2011;121:1187-93.
3. Rzepakowska A, Osuch-Wójcikiewicz E, Kulesza A, Bruzgielewicz A, Niemczyk K. [Paragangliomas of the neck: A 10-year experience of the department of otolaryngology of Warsaw medical university]. *Otolaryngol Pol* 2010;64:65-72.
4. Sanders KW, Abreo F, Rivera E, Stucker FJ, Nathan CO. A diagnostic and therapeutic approach to paragangliomas of the larynx. *Arch Otolaryngol Head Neck Surg* 2001;127:565-9.

Beigi and Eshaghian: Superior laryngeal paraganglioma

5. Kaytaz A, Karaman E, Edizer DT, Hacıyev Y, Öz B. Laryngeal paraganglioma: Report of an unusual entity. *Ear Nose Throat J* 2010;89:550-2.
6. Wang SH, Chiu KM, Cheng PW. Bilateral carotid body paragangliomas. *CMAJ* 2011;183:E606.
7. Ghoreishi M, Akbar-Beigi A, Tahery D, Sehat S. Fever as the main presenting symptom of a carotid body tumor. *Arch Iran Med* 2008;11:214-7.
8. Ma D, Liu M, Yang H, Ma X, Zhang C. Diagnosis and surgical treatment of carotid body tumor: A report of 18 cases. *J Cardiovasc Dis Res* 2010;1:122-4.
9. Rubio V, Tamarit JM, Baviera N, Estrems P, López C, Estellés E, et al. [Laryngeal paraganglioma: Diagnosis and treatment. A propos of a case]. *Acta Otorrinolaringol Esp* 2007;58:333-4.
10. Buiret G, Merrot O, Le Breton F, Landry G, Pignat JC. Supraglottic laryngeal paraganglioma. *Eur Ann Otorhinolaryngol Head Neck Dis* 2010;127:117-9.
11. Takumida M, Taira T, Suzuki M, Yajin K, Harada Y. Neurilemmoma of the larynx: A case report. *J Laryngol Otol* 1986;100:847-50.
12. Papacharalampous GX, Korres S, Tzagaroulakis M, Segas I, Ferekidis E. Paraganglioma of the larynx: A case report. *Med Sci Monit* 2007;13:CS145-8.
13. Capolunghi B, Bertolini G, Grillodella Berta L, Tinelli N, Cascio F, Bertolotti F. Laryngeal paraganglioma: An endoscopic diode-laser-assisted surgical approach: A case report. *B-ENT* 2005;1:97-100.

Source of Support: Nil, **Conflict of Interest:** None declared.